

Goat Anti-ANILLIN / Scraps (internal) Antibody
Peptide-affinity purified goat antibody
Catalog # AF1062b

Specification

Goat Anti-ANILLIN / Scraps (internal) Antibody - Product Information

Application	WB, Pep-ELISA
Primary Accession	Q9NOW6
Other Accession	NP_061155 , 54443 , 68743 (mouse) , 307056 (rat)
Reactivity	Human, Mouse
Predicted	Rat, Dog
Host	Goat
Clonality	Polyclonal
Concentration	100ug/200ul
Isotype	IgG
Calculated MW	124199

Goat Anti-ANILLIN / Scraps (internal) Antibody - Additional Information

Gene ID 54443

Other Names

Actin-binding protein anillin, ANLN

Dilution

WB~~1:1000
Pep-ELISA~~N/A

Format

0.5 mg IgG/ml in Tris saline (20mM Tris pH7.3, 150mM NaCl), 0.02% sodium azide, with 0.5% bovine serum albumin

Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

Goat Anti-ANILLIN / Scraps (internal) Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

Goat Anti-ANILLIN / Scraps (internal) Antibody - Protein Information

Name ANLN

Function

Required for cytokinesis (PubMed:<http://www.uniprot.org/citations/16040610>)

target="_blank">>16040610). Essential for the structural integrity of the cleavage furrow and for completion of cleavage furrow ingression. Plays a role in bleb assembly during metaphase and anaphase of mitosis (PubMed:23870127). May play a significant role in podocyte cell migration (PubMed:24676636).

Cellular Location

Nucleus. Cytoplasm, cytoskeleton. Cytoplasm, cell cortex. Cell projection, bleb. Note=Mainly found in the nucleus during interphase. Colocalizes with cortical F-actin upon nuclear envelope breakdown in mitosis and subsequently concentrates in the area of the prospective contractile ring in anaphase. This pattern persists until telophase, when the protein becomes concentrated in the midbody

Tissue Location

Ubiquitously expressed. Present at highest levels in the brain, at high levels in the placenta and testis, at intermediate levels in the intestine, ovary, skeletal muscle and thymus and at lower levels in heart, kidney, liver, lung, pancreas, prostate and spleen. In the kidney, it is widely expressed in tubules, but sparsely expressed in the glomerulus (PubMed:24676636). Expression is significantly increased in renal biopsy specimens from idiopathic FSGS (PubMed:24676636). Overexpressed in many tumor types including breast, colorectal, endometrial, hepatic, kidney, lung, ovarian and pancreatic tumors.

Goat Anti-ANILLIN / Scraps (internal) Antibody - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)


Goat Anti-ANILLIN / Scraps (internal) Antibody - Images



AF1062b (1 µg/ml) staining of Mouse Brain lysate (35 µg protein in RIPA buffer). Primary incubation was 1 hour. Detected by chemiluminescence.



EB07300 Immunofluorescence analysis of paraformaldehyde fixed NIH3T3 cells, permeabilized with 0.15% Triton. Primary incubation 1hr (10ug/ml) followed by Alexa Fluor 488 secondary antibody (2ug/ml), showing nuclear and cytoplasmic staining. The nuclear stain is DAPI (blue). Negative control: Unimmunized goat IgG (10ug/ml) followed by Alexa Fluor 488 secondary antibody (2ug/ml).



EB07300 Immunofluorescence analysis of paraformaldehyde fixed Neuro-2a cells, permeabilized with 0.15% Triton. Primary incubation 1hr (10ug/ml) followed by Alexa Fluor 488 secondary antibody (2ug/ml), showing nuclear staining. The nuclear stain is DAPI (blue). Negative control: Unimmunized goat IgG (10ug/ml) followed by Alexa Fluor 488 secondary antibody (2ug/ml).

Goat Anti-ANILLIN / Scraps (internal) Antibody - References

NBL1 and anillin (ANLN) genes over-expression in pancreatic carcinoma. Olakowski M, et al. *Folia Histochem Cytobiol*, 2009. PMID 19995712.

Anillin is a scaffold protein that links RhoA, actin, and myosin during cytokinesis. Piekny AJ, et al. *Curr Biol*, 2008 Jan 8. PMID 18158243.

Systematic analysis of the protein interaction network for the human transcription machinery reveals the identity of the 7SK capping enzyme. Jeronimo C, et al. *Mol Cell*, 2007 Jul 20. PMID 17643375.

Global, *in vivo*, and site-specific phosphorylation dynamics in signaling networks. Olsen JV, et al. *Cell*, 2006 Nov 3. PMID 17081983.

A probability-based approach for high-throughput protein phosphorylation analysis and site localization. Beausoleil SA, et al. Nat Biotechnol, 2006 Oct. PMID 16964243.