

Goat Anti-CAMK1D Antibody
Peptide-affinity purified goat antibody
Catalog # AF1183a**Specification**

Goat Anti-CAMK1D Antibody - Product Information

Application	WB, E
Primary Accession	Q8IU85
Other Accession	NP_065130 , 57118
Reactivity	Human
Host	Goat
Clonality	Polyclonal
Concentration	100ug/200ul
Isotype	IgG
Calculated MW	42914

Goat Anti-CAMK1D Antibody - Additional Information**Gene ID** 57118**Other Names**

Calcium/calmodulin-dependent protein kinase type 1D, 2.7.11.17, CaM kinase I delta, CaM kinase ID, CaM-KI delta, CaMKI delta, CaMKID, CaMKI-like protein kinase, CKLiK, CAMK1D, CAMKID

Dilution

WB~~1:1000

E~~N/A

Format

0.5 mg IgG/ml in Tris saline (20mM Tris pH7.3, 150mM NaCl), 0.02% sodium azide, with 0.5% bovine serum albumin

Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

Goat Anti-CAMK1D Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

Goat Anti-CAMK1D Antibody - Protein Information**Name** CAMK1D**Synonyms** CAMKID**Function**

Calcium/calmodulin-dependent protein kinase that operates in the calcium-triggered CaMKK-CaMK1 signaling cascade and, upon calcium influx, activates CREB-dependent gene transcription, regulates calcium-mediated granulocyte function and respiratory burst and promotes basal dendritic growth of hippocampal neurons. In neutrophil cells, required for cytokine-induced proliferative responses and activation of the respiratory burst. Activates the transcription factor CREB1 in hippocampal neuron nuclei. May play a role in apoptosis of erythroleukemia cells. In vitro, phosphorylates transcription factor CREM isoform Beta.

Cellular Location

Cytoplasm. Nucleus. Note=Predominantly cytoplasmic. Nuclear localization increases upon activation by KCl treatment in hippocampal neurons

Tissue Location

Widely expressed. Highly and mostly expressed in polymorphonuclear leukocytes (neutrophilic and eosinophilic granulocytes) while little or no expression is observed in monocytes and lymphocytes.

Goat Anti-CAMK1D Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Goat Anti-CAMK1D Antibody - Images



AF1183a (0.5 µg/ml) staining of Human Liver lysate (35 µg protein in RIPA buffer). Primary incubation was 1 hour. Detected by chemiluminescence.

Goat Anti-CAMK1D Antibody - Background

This gene encodes a member of the Ca²⁺/calmodulin-dependent protein kinase 1 subfamily of serine/threonine kinases. The encoded protein may be involved in the regulation of granulocyte function through the chemokine signal transduction pathway. Alternatively spliced transcript variants encoding different isoforms of this gene have been described.

Goat Anti-CAMK1D Antibody - References

An approach based on a genome-wide association study reveals candidate loci for narcolepsy. Shimada M, et al. Hum Genet, 2010 Oct. PMID 20677014.

Evaluating the discriminative power of multi-trait genetic risk scores for type 2 diabetes in a northern Swedish population. Fontaine-Bisson B, et al. Diabetologia, 2010 Oct. PMID 20571754.

Common variants associated with breast cancer in genome-wide association studies are modifiers of breast cancer risk in BRCA1 and BRCA2 mutation carriers. Wang X, et al. Hum Mol Genet, 2010 Jul 15. PMID 20418484.

Personalized smoking cessation: interactions between nicotine dose, dependence and quit-success genotype score. Rose JE, et al. Mol Med, 2010 Jul-Aug. PMID 20379614.

Investigation of type 2 diabetes risk alleles support CDKN2A/B, CDKAL1, and TCF7L2 as susceptibility genes in a Han Chinese cohort. Wen J, et al. PLoS One, 2010 Feb 10. PMID 20161779.