10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

Goat Anti-CAMK1D Antibody

Peptide-affinity purified goat antibody

Catalog # AF1183a

Specification

Goat Anti-CAMK1D Antibody - Product Information

Application	**WB**
Primary Accession	Q8IU85
Other Accession	NP_065130, 57118
Reactivity	**Human**
Host	**Goat**
Clonality	**Polyclonal**
Concentration	**100ug/200ul**
Isotype	**IgG**
Calculated MW	**42914**

Goat Anti-CAMK1D Antibody - Additional Information

Gene ID 57118

Other Names
Calcium/calmodulin-dependent protein kinase type 1D, 2.7.11.17, CaM kinase I delta, CaM kinase ID, CaM-KI delta, CaMKI delta, CaMKID, CaMKI-like protein kinase, CKLiK, CAMK1D, CAMKID

Format
0.5 mg IgG/ml in Tris saline (20mM Tris pH7.3, 150mM NaCl), 0.02% sodium azide, with 0.5% bovine serum albumin

Storage
Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions
Goat Anti-CAMK1D Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

Goat Anti-CAMK1D Antibody - Protein Information

Name CAMK1D

Synonyms CAMKID

Function
Calcium/calmodulin-dependent protein kinase that operates in the calcium-triggered CaMKK-CaMK1 signaling cascade and, upon calcium influx, activates CREB-dependent gene transcription, regulates calcium- mediated granulocyte function and respiratory burst and promotes basal dendritic growth of hippocampal neurons. In neutrophil cells, required for

cytokine-induced proliferative responses and activation of the respiratory burst. Activates the transcription factor CREB1 in hippocampal neuron nuclei. May play a role in apoptosis of erythroleukemia cells. In vitro, phosphorylates transcription factor CREM isoform Beta.

Cellular Location
Cytoplasm. Nucleus. Note=Predominantly cytoplasmic. Nuclear localization increases upon activation by KCl treatment in hippocampal neurons

Tissue Location
Widely expressed. Highly and mostly expressed in polymorphonuclear leukocytes (neutrophilic and eosinophilic granulocytes) while little or no expression is observed in monocytes and lymphocytes.

Goat Anti-CAMK1D Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

Goat Anti-CAMK1D Antibody - Images



AF1183a (0.5 µg/ml) staining of Human Liver lysate (35 µg protein in RIPA buffer). Primary incubation was 1 hour. Detected by chemiluminescence.

Goat Anti-CAMK1D Antibody - Background

This gene encodes a member of the Ca2+/calmodulin-dependent protein kinase 1 subfamily of serine/threonine kinases. The encoded protein may be involved in the regulation of granulocyte function through the chemokine signal transduction pathway. Alternatively spliced transcript variants encoding different isoforms of this gene have been described.

Goat Anti-CAMK1D Antibody - References

An approach based on a genome-wide association study reveals candidate loci for narcolepsy. Shimada M, et al. Hum Genet, 2010 Oct. PMID 20677014.
Evaluating the discriminative power of multi-trait genetic risk scores for type 2 diabetes in a

northern Swedish population. Fontaine-Bisson B, et al. Diabetologia, 2010 Oct. PMID 20571754.
Common variants associated with breast cancer in genome-wide association studies are modifiers of breast cancer risk in BRCA1 and BRCA2 mutation carriers. Wang X, et al. Hum Mol Genet, 2010 Jul 15. PMID 20418484.
Personalized smoking cessation: interactions between nicotine dose, dependence and quit-success genotype score. Rose JE, et al. Mol Med, 2010 Jul-Aug. PMID 20379614.
Investigation of type 2 diabetes risk alleles support CDKN2A/B, CDKAL1, and TCF7L2 as susceptibility genes in a Han Chinese cohort. Wen J, et al. PLoS One, 2010 Feb 10. PMID 20161779.