

**Goat Anti-Cyclophilin A / PPIA Antibody**  
**Peptide-affinity purified goat antibody**  
**Catalog # AF1290a****Specification**

---

**Goat Anti-Cyclophilin A / PPIA Antibody - Product Information**

Application	WB, E
Primary Accession	<a href="#">P62937</a>
Other Accession	<a href="#">NP_066953</a> , <a href="#">5478</a>
Reactivity	Human
Predicted	Mouse, Rat, Pig, Dog
Host	Goat
Clonality	Polyclonal
Concentration	100ug/200ul
Isotype	IgG
Calculated MW	18012

**Goat Anti-Cyclophilin A / PPIA Antibody - Additional Information****Gene ID** 5478**Other Names**

Peptidyl-prolyl cis-trans isomerase A, PPIase A, 5.2.1.8, Cyclophilin A, Cyclosporin A-binding protein, Rotamase A, Peptidyl-prolyl cis-trans isomerase A, N-terminally processed, PPIA, CYPA

**Dilution**

WB~~1:1000

E~~N/A

**Format**

0.5 mg IgG/ml in Tris saline (20mM Tris pH7.3, 150mM NaCl), 0.02% sodium azide, with 0.5% bovine serum albumin

**Storage**

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

**Precautions**

Goat Anti-Cyclophilin A / PPIA Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

**Goat Anti-Cyclophilin A / PPIA Antibody - Protein Information****Name** PPIA**Synonyms** CYPA

## Function

Catalyzes the cis-trans isomerization of proline imidic peptide bonds in oligopeptides (PubMed:<a href="http://www.uniprot.org/citations/2001362" target="\_blank">2001362</a>, PubMed:<a href="http://www.uniprot.org/citations/20676357" target="\_blank">20676357</a>, PubMed:<a href="http://www.uniprot.org/citations/21245143" target="\_blank">21245143</a>, PubMed:<a href="http://www.uniprot.org/citations/21593166" target="\_blank">21593166</a>, PubMed:<a href="http://www.uniprot.org/citations/25678563" target="\_blank">25678563</a>). Exerts a strong chemotactic effect on leukocytes partly through activation of one of its membrane receptors BSG/CD147, initiating a signaling cascade that culminates in MAPK/ERK activation (PubMed:<a href="http://www.uniprot.org/citations/11943775" target="\_blank">11943775</a>, PubMed:<a href="http://www.uniprot.org/citations/21245143" target="\_blank">21245143</a>). Activates endothelial cells (ECs) in a pro-inflammatory manner by stimulating activation of NF-kappa-B and ERK, JNK and p38 MAP-kinases and by inducing expression of adhesion molecules including SELE and VCAM1 (PubMed:<a href="http://www.uniprot.org/citations/15130913" target="\_blank">15130913</a>). Induces apoptosis in ECs by promoting the FOXO1-dependent expression of CCL2 and BCL2L1 which are involved in EC chemotaxis and apoptosis (PubMed:<a href="http://www.uniprot.org/citations/31063815" target="\_blank">31063815</a>). In response to oxidative stress, initiates proapoptotic and antiapoptotic signaling in ECs via activation of NF-kappa-B and AKT1 and up-regulation of antiapoptotic protein BCL2 (PubMed:<a href="http://www.uniprot.org/citations/23180369" target="\_blank">23180369</a>). Negatively regulates MAP3K5/ASK1 kinase activity, autophosphorylation and oxidative stress-induced apoptosis mediated by MAP3K5/ASK1 (PubMed:<a href="http://www.uniprot.org/citations/26095851" target="\_blank">26095851</a>). Necessary for the assembly of TARDBP in heterogeneous nuclear ribonucleoprotein (hnRNP) complexes and regulates TARDBP binding to RNA UG repeats and TARDBP-dependent expression of HDAC6, ATG7 and VCP which are involved in clearance of protein aggregates (PubMed:<a href="http://www.uniprot.org/citations/25678563" target="\_blank">25678563</a>). Plays an important role in platelet activation and aggregation (By similarity). Regulates calcium mobilization and integrin ITGA2B:ITGB3 bidirectional signaling via increased ROS production as well as by facilitating the interaction between integrin and the cell cytoskeleton (By similarity). Binds heparan sulfate glycosaminoglycans (PubMed:<a href="http://www.uniprot.org/citations/11943775" target="\_blank">11943775</a>). Inhibits replication of influenza A virus (IAV) (PubMed:<a href="http://www.uniprot.org/citations/19207730" target="\_blank">19207730</a>). Inhibits ITCH/AIP4-mediated ubiquitination of matrix protein 1 (M1) of IAV by impairing the interaction of ITCH/AIP4 with M1, followed by the suppression of the nuclear export of M1, and finally reduction of the replication of IAV (PubMed:<a href="http://www.uniprot.org/citations/22347431" target="\_blank">22347431</a>, PubMed:<a href="http://www.uniprot.org/citations/30328013" target="\_blank">30328013</a>).

## Cellular Location

Cytoplasm. Secreted. Nucleus Note=Secretion occurs in response to oxidative stress in vascular smooth muscle through a vesicular secretory pathway that includes Rho GTPase signaling, actin remodeling, and myosin II activation

## Goat Anti-Cyclophilin A / PPIA Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)

- [Flow Cytometry](#)
- [Cell Culture](#)

### Goat Anti-Cyclophilin A / PPIA Antibody - Images



AF1290a (0.1 µg/ml) staining of Human Lymph lysate (35 µg protein in RIPA buffer). Primary incubation was 1 hour. Detected by chemiluminescence.

### Goat Anti-Cyclophilin A / PPIA Antibody - Background

This gene encodes a member of the peptidyl-prolyl cis-trans isomerase (PPIase) family. PPIases catalyze the cis-trans isomerization of proline imidic peptide bonds in oligopeptides and accelerate the folding of proteins. The encoded protein is a cyclosporin binding-protein and may play a role in cyclosporin A-mediated immunosuppression. The protein can also interact with several HIV proteins, including p55 gag, Vpr, and capsid protein, and has been shown to be necessary for the formation of infectious HIV virions. Multiple pseudogenes that map to different chromosomes have been reported.

### Goat Anti-Cyclophilin A / PPIA Antibody - References

Proteome analysis of the thalamus and cerebrospinal fluid reveals glycolysis dysfunction and potential biomarkers candidates for schizophrenia. Martins-de-Souza D, et al. J Psychiatr Res, 2010 May 14. PMID 20471030.

Sex-specific proteome differences in the anterior cingulate cortex of schizophrenia.

Martins-de-Souza D, et al. J Psychiatr Res, 2010 Apr 8. PMID 20381070.

SERCA2b activity is regulated by cyclophilins in human platelets. Rosado JA, et al. Arterioscler Thromb Vasc Biol, 2010 Mar. PMID 20139366.

Cyclophilin A-independent recruitment of NS5A and NS5B into hepatitis C virus replication complexes. Chatterji U, et al. J Gen Virol, 2010 May. PMID 20107018.

Chaperone activity of Cyp18 through hydrophobic condensation that enables rescue of transient misfolded molten globule intermediates. Moparthi SB, et al. Biochemistry, 2010 Feb 16. PMID 20070121.