

**Goat Anti-DLG4 / PSD95 Antibody**  
**Peptide-affinity purified goat antibody**  
**Catalog # AF1329a****Specification**

---

**Goat Anti-DLG4 / PSD95 Antibody - Product Information**

Application	WB, E
Primary Accession	<a href="#">P78352</a>
Other Accession	<a href="#">NP_001122299</a> , <a href="#">1742</a> , <a href="#">13385 (mouse)</a> , <a href="#">29495 (rat)</a>
Reactivity	Rat
Predicted	Human, Mouse, Dog
Host	Goat
Clonality	Polyclonal
Concentration	100ug/200ul
Isotype	IgG
Calculated MW	80495

**Goat Anti-DLG4 / PSD95 Antibody - Additional Information****Gene ID** 1742**Other Names**

Disks large homolog 4, Postsynaptic density protein 95, PSD-95, Synapse-associated protein 90, SAP-90, SAP90, DLG4, PSD95

**Dilution**

WB~~1:1000

E~~N/A

**Format**

0.5 mg IgG/ml in Tris saline (20mM Tris pH7.3, 150mM NaCl), 0.02% sodium azide, with 0.5% bovine serum albumin

**Storage**

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

**Precautions**

Goat Anti-DLG4 / PSD95 Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

**Goat Anti-DLG4 / PSD95 Antibody - Protein Information****Name** DLG4 ([HGNC:2903](#))**Synonyms** PSD95

**Function**

Postsynaptic scaffolding protein that plays a critical role in synaptogenesis and synaptic plasticity by providing a platform for the postsynaptic clustering of crucial synaptic proteins. Interacts with the cytoplasmic tail of NMDA receptor subunits and shaker-type potassium channels. Required for synaptic plasticity associated with NMDA receptor signaling. Overexpression or depletion of DLG4 changes the ratio of excitatory to inhibitory synapses in hippocampal neurons. May reduce the amplitude of ASIC3 acid-evoked currents by retaining the channel intracellularly. May regulate the intracellular trafficking of ADR1B. Also regulates AMPA-type glutamate receptor (AMPA) immobilization at postsynaptic density keeping the channels in an activated state in the presence of glutamate and preventing synaptic depression (By similarity). Under basal conditions, cooperates with FYN to stabilize palmitoyltransferase ZDHHC5 at the synaptic membrane through FYN-mediated phosphorylation of ZDHHC5 and its subsequent inhibition of association with endocytic proteins (PubMed:<a href="http://www.uniprot.org/citations/26334723" target="\_blank">26334723</a>).

**Cellular Location**

Cell membrane; Lipid-anchor; Cytoplasmic side. Postsynaptic density {ECO:0000250|UniProtKB:P31016}. Synapse Cytoplasm {ECO:0000250|UniProtKB:P31016}. Cell projection, axon {ECO:0000250|UniProtKB:P31016}. Cell projection, dendritic spine {ECO:0000250|UniProtKB:P31016}. Cell projection, dendrite {ECO:0000250|UniProtKB:P31016}. Presynapse {ECO:0000250|UniProtKB:P31016}. Note=High levels in postsynaptic density of neurons in the forebrain. Also in presynaptic region of inhibitory synapses formed by cerebellar basket cells on axon hillocks of Purkinje cells. Suppression of neuronal activity induces synaptic accumulation and clustering of DLG4. {ECO:0000250|UniProtKB:P31016}

**Tissue Location**


Brain.

**Goat Anti-DLG4 / PSD95 Antibody - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

**Goat Anti-DLG4 / PSD95 Antibody - Images**



AF1329a (0.3 µg/ml) staining of Rat Brain lysate (35 µg protein in RIPA buffer). Primary incubation was 1 hour. Detected by chemiluminescence.

#### **Goat Anti-DLG4 / PSD95 Antibody - Background**

This gene encodes a member of the membrane-associated guanylate kinase (MAGUK) family. It heteromultimerizes with another MAGUK protein, DLG2, and is recruited into NMDA receptor and potassium channel clusters. These two MAGUK proteins may interact at postsynaptic sites to form a multimeric scaffold for the clustering of receptors, ion channels, and associated signaling proteins. Multiple transcript variants encoding different isoforms have been found for this gene.

#### **Goat Anti-DLG4 / PSD95 Antibody - References**

Variation at the NFATC2 Locus Increases the Risk of Thiazolidinedione-Induced Edema in the Diabetes REduction Assessment with ramipril and rosiglitazone Medication (DREAM) Study. Bailey SD, et al. Diabetes Care, 2010 Jul 13. PMID 20628086.

Human variation in alcohol response is influenced by variation in neuronal signaling genes. Joslyn G, et al. Alcohol Clin Exp Res, 2010 May. PMID 20201926.

Gene-centric association signals for lipids and apolipoproteins identified via the HumanCVD BeadChip. Talmud PJ, et al. Am J Hum Genet, 2009 Nov. PMID 19913121.

Decreased levels of PSD95 and two associated proteins and increased levels of BCL2 and caspase 3 in hippocampus from subjects with amnesic mild cognitive impairment: Insights into their potential roles for loss of synapses and memory, accumulation of Abeta, and neurodegeneration in a prodromal stage of Alzheimer's disease. Sultana R, et al. J Neurosci Res, 2010 Feb 15. PMID 19774677.

Interaction of postsynaptic density protein-95 with NMDA receptors influences excitotoxicity in the yeast artificial chromosome mouse model of Huntington's disease. Fan J, et al. J Neurosci, 2009 Sep 2. PMID 19726651.