

Goat Anti-DLL1 Antibody
Peptide-affinity purified goat antibody
Catalog # AF1330a**Specification**

Goat Anti-DLL1 Antibody - Product Information

Application	WB, E
Primary Accession	O00548
Other Accession	NP_005609 , 28514 , 13388 (mouse) , 84010 (rat)
Reactivity	Mouse, Rat
Predicted	Human
Host	Goat
Clonality	Polyclonal
Concentration	100ug/200ul
Isotype	IgG
Calculated MW	78056

Goat Anti-DLL1 Antibody - Additional Information**Gene ID** 28514**Other Names**

Delta-like protein 1, Drosophila Delta homolog 1, Delta1, H-Delta-1, DLL1

Dilution

WB~~1:1000

E~~N/A

Format

0.5 mg IgG/ml in Tris saline (20mM Tris pH7.3, 150mM NaCl), 0.02% sodium azide, with 0.5% bovine serum albumin

Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

Goat Anti-DLL1 Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

Goat Anti-DLL1 Antibody - Protein Information**Name** DLL1 ([HGNC:2908](#))**Function**

Transmembrane ligand protein of NOTCH1, NOTCH2 and NOTCH3 receptors that binds the extracellular domain (ECD) of Notch receptor in a cis and trans fashion manner (PubMed:<a

<http://www.uniprot.org/citations/11006133> (11006133). Following transinteraction, ligand cells produce mechanical force that depends of a clathrin-mediated endocytosis, requiring ligand ubiquitination, EPN1 interaction, and actin polymerisation; these events promote Notch receptor extracellular domain (NECD) transendocytosis and triggers Notch signaling through induction of cleavage, hyperphosphorylation, and nuclear accumulation of the intracellular domain of Notch receptors (NICD) (By similarity). Is required for embryonic development and maintenance of adult stem cells in many different tissues and immune systems; the DLL1-induced Notch signaling is mediated through an intercellular communication that regulates cell lineage, cell specification, cell patterning and morphogenesis through effects on differentiation and proliferation (PubMed: <http://www.uniprot.org/citations/11581320> (11581320)). Plays a role in brain development at different level, namely by regulating neuronal differentiation of neural precursor cells via cell-cell interaction, most likely through the lateral inhibitory system in an endogenous level dependent-manner. During neocortex development, DLL1-Notch signaling transmission is mediated by dynamic interactions between intermediate neurogenic progenitors and radial glia; the cell-cell interactions are mediated via dynamic and transient elongation processes, likely to reactivate/maintain Notch activity in neighboring progenitors, and coordinate progenitor cell division and differentiation across radial and zonal boundaries. During cerebellar development, regulates Bergmann glial monolayer formation and its morphological maturation through a Notch signaling pathway. At the retina and spinal cord level, regulates neurogenesis by preventing the premature differentiation of neural progenitors and also by maintaining progenitors in spinal cord through Notch signaling pathway. Also controls neurogenesis of the neural tube in a progenitor domain-specific fashion along the dorsoventral axis. Maintains quiescence of neural stem cells and plays a role as a fate determinant that segregates asymmetrically to one daughter cell during neural stem cells mitosis, resulting in neuronal differentiation in DLL1-inheriting cell. Plays a role in immune system development, namely the development of all T-cells and marginal zone (MZ) B-cells (By similarity). Blocks the differentiation of progenitor cells into the B-cell lineage while promoting the emergence of a population of cells with the characteristics of a T-cell/NK-cell precursor (PubMed: <http://www.uniprot.org/citations/11581320> (11581320)). Also plays a role during muscle development. During early development, inhibits myoblasts differentiation from the medial dermomyotomal lip and later regulates progenitor cell differentiation. Directly modulates cell adhesion and basal lamina formation in satellite cells through Notch signaling. Maintains myogenic progenitors pool by suppressing differentiation through down-regulation of MYOD1 and is required for satellite cell homing and PAX7 expression. During craniofacial and trunk myogenesis suppresses differentiation of cranial mesoderm-derived and somite-derived muscle via MYOD1 regulation but in cranial mesoderm-derived progenitors, is neither required for satellite cell homing nor for PAX7 expression. Also plays a role during pancreatic cell development. During type B pancreatic cell development, may be involved in the initiation of proximodistal patterning in the early pancreatic epithelium. Stimulates multipotent pancreatic progenitor cells proliferation and pancreatic growth by maintaining HES1 expression and PTF1A protein levels. During fetal stages of development, is required to maintain arterial identity and the responsiveness of arterial endothelial cells for VEGFA through regulation of KDR activation and NRP1 expression. Controls sprouting angiogenesis and subsequent vertical branch formation through regulation on tip cell differentiation. Negatively regulates goblet cell differentiation in intestine and controls secretory fat commitment through lateral inhibition in small intestine. Plays a role during inner ear development; negatively regulates auditory hair cell differentiation. Plays a role during nephron development through Notch signaling pathway. Regulates growth, blood pressure and energy homeostasis (By similarity).

Cellular Location

Apical cell membrane {ECO:0000250|UniProtKB:Q61483}; Single-pass type I membrane protein {ECO:0000250|UniProtKB:Q61483}. Cell junction, adherens junction {ECO:0000250|UniProtKB:Q61483}. Membrane raft {ECO:0000250|UniProtKB:Q61483}. Note=Distributed around adherens junction in the apical endfeet through interactions with MAGI1 {ECO:0000250|UniProtKB:Q61483}

Tissue Location

Expressed in heart and pancreas, with lower expression in brain and muscle and almost no expression in placenta, lung, liver and kidney

Goat Anti-DLL1 Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Goat Anti-DLL1 Antibody - Images



AF1330a (0.3 µg/ml) staining of Rat Heart lysate (35 µg protein in RIPA buffer). Primary incubation was 1 hour. Detected by chemiluminescence.

Goat Anti-DLL1 Antibody - Background

DLL1 is a human homolog of the Notch Delta ligand and is a member of the delta/serrate/jagged family. It plays a role in mediating cell fate decisions during hematopoiesis. It may play a role in cell-to-cell communication.

Goat Anti-DLL1 Antibody - References

Squamous odontogenic tumor of the mandible: a case report demonstrating immunoexpression of Notch1, 3, 4, Jagged1 and Delta1. Siar CH, et al. Eur J Med Res, 2010 Apr 8. PMID 20554499.
Cis-interactions between Notch and Delta generate mutually exclusive signalling states. Sprinzak D, et al. Nature, 2010 May 6. PMID 20418862.
Notch1 induces enhanced expression of Delta-like-1 in the U251MG glioma cell line. Qian CF, et al. Int J Mol Med, 2009 Oct. PMID 19724883.
Genetic variation in healthy oldest-old. Halaschek-Wiener J, et al. PLoS One, 2009 Aug 14. PMID 19680556.
Aberrant expression of Notch signaling molecules in patients with immune thrombocytopenic purpura. Ma D, et al. Ann Hematol, 2010 Feb. PMID 19603167.