

Goat Anti-EAT2 phospho (Y127) Antibody
Peptide-affinity purified goat antibody
Catalog # AF1351a**Specification**

Goat Anti-EAT2 phospho (Y127) Antibody - Product Information

Application	WB, E
Primary Accession	O14796
Other Accession	NP_444512 , 117157
Reactivity	Human
Host	Goat
Clonality	Polyclonal
Concentration	100ug/200ul
Isotype	IgG
Calculated MW	15297

Goat Anti-EAT2 phospho (Y127) Antibody - Additional Information**Gene ID** 117157**Other Names**

SH2 domain-containing protein 1B, EWS/FLI1-activated transcript 2, EAT-2, SH2D1B, EAT2

Format

0.5 mg IgG/ml in Tris saline (20mM Tris pH7.3, 150mM NaCl), 0.02% sodium azide, with 0.5% bovine serum albumin

Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

Goat Anti-EAT2 phospho (Y127) Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

Goat Anti-EAT2 phospho (Y127) Antibody - Protein Information**Name** SH2D1B**Synonyms** EAT2**Function**

Cytoplasmic adapter regulating receptors of the signaling lymphocytic activation molecule (SLAM) family such as CD84, SLAMF1, LY9 and CD244 (PubMed:11689425). In SLAM signaling seems to cooperate with SH2D1A/SAP. Plays a role in regulation of effector functions of natural killer (NK) cells by controlling signal transduction through CD244/2B4 without effecting its

tyrosine phosphorylation; downstream signaling involves PLCG1 and ERK activation (PubMed:24687958). Activation of SLAMF7-mediated NK cell function does not effect receptor tyrosine phosphorylation but distal signaling (By similarity). In the context of NK cell-mediated cytotoxicity does not enhance conjugate formation with target cells but stimulates polarization of the microtubule-organizing center and cytotoxic granules toward the NK cell synapse (PubMed:24687958). Negatively regulates CD40-induced cytokine production in dendritic cells downstream of SLAM family receptors probably by inducing activation of the PI3K pathway to inhibit p38 MAPK and JNK activation (By similarity).

Goat Anti-EAT2 phospho (Y127) Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Goat Anti-EAT2 phospho (Y127) Antibody - Images

Goat Anti-EAT2 phospho (Y127) Antibody - Background

By binding phosphotyrosines through its free SRC (MIM 190090) homology-2 (SH2) domain, EAT2 regulates signal transduction through receptors expressed on the surface of antigen-presenting cells (Morra et al., 2001 [PubMed 11689425]).

Goat Anti-EAT2 phospho (Y127) Antibody - References

Molecular analysis of NTB-A signaling: a role for EAT-2 in NTB-A-mediated activation of human NK cells. Eissmann P, et al. J Immunol, 2006 Sep 1. PMID 16920955.
The DNA sequence and biological annotation of human chromosome 1. Gregory SG, et al. Nature, 2006 May 18. PMID 16710414.
The status, quality, and expansion of the NIH full-length cDNA project: the Mammalian Gene Collection (MGC). Gerhard DS, et al. Genome Res, 2004 Oct. PMID 15489334.
Complete sequencing and characterization of 21,243 full-length human cDNAs. Ota T, et al. Nat Genet, 2004 Jan. PMID 14702039.
The SLAM family of immune-cell receptors. Veillette A, et al. Curr Opin Immunol, 2003 Jun. PMID 12787752.