

Goat Anti-FANCG / XRCC9 Antibody
Peptide-affinity purified goat antibody
Catalog # AF1399a**Specification**

Goat Anti-FANCG / XRCC9 Antibody - Product Information

Application	WB
Primary Accession	O15287
Other Accession	NP_004620 , 2189
Reactivity	Human
Predicted	Pig, Dog
Host	Goat
Clonality	Polyclonal
Concentration	100ug/200ul
Isotype	IgG
Calculated MW	68554

Goat Anti-FANCG / XRCC9 Antibody - Additional Information**Gene ID** 2189**Other Names**

Fanconi anemia group G protein, Protein FACG, DNA repair protein XRCC9, FANCG, XRCC9

Format

0.5 mg IgG/ml in Tris saline (20mM Tris pH7.3, 150mM NaCl), 0.02% sodium azide, with 0.5% bovine serum albumin

Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

Goat Anti-FANCG / XRCC9 Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

Goat Anti-FANCG / XRCC9 Antibody - Protein Information**Name** FANCG**Synonyms** XRCC9**Function**

DNA repair protein that may operate in a postreplication repair or a cell cycle checkpoint function. May be implicated in interstrand DNA cross-link repair and in the maintenance of normal chromosome stability. Candidate tumor suppressor gene.

Cellular Location

Nucleus. Cytoplasm. Note=The major form is nuclear. The minor form is cytoplasmic

Tissue Location

Highly expressed in testis and thymus. Found in lymphoblasts


Goat Anti-FANCG / XRCC9 Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Goat Anti-FANCG / XRCC9 Antibody - Images

AF1399a (0.5 µg/ml) staining of HeLa cell lysate (35 µg protein in RIPA buffer). Primary incubation was 1 hour. Detected by chemiluminescence.



HEK293 overexpressing FANCG (RC202443) and probed with AF1399a (mock transfection in first lane), tested by Origene.

Goat Anti-FANCG / XRCC9 Antibody - Background

The Fanconi anemia complementation group (FANC) currently includes FANCA, FANCB, FANCC, FANCD1 (also called BRCA2), FANCD2, FANCE, FANCF, FANCG, FANCI, FANCI (also called BRIP1), FANCL, FANCM and FANCN (also called PALB2). The previously defined group FANCH is the same as FANCA. Fanconi anemia is a genetically heterogeneous recessive disorder characterized by cytogenetic instability, hypersensitivity to DNA crosslinking agents, increased chromosomal breakage, and defective DNA repair. The members of the Fanconi anemia complementation group do not share sequence similarity; they are related by their assembly into a common nuclear protein complex. This gene encodes the protein for complementation group G.

Goat Anti-FANCG / XRCC9 Antibody - References

Variation at the NFATC2 Locus Increases the Risk of Thiazolinedione-Induced Edema in the Diabetes REduction Assessment with ramipril and rosiglitazone Medication (DREAM) Study. Bailey SD, et al. Diabetes Care, 2010 Jul 13. PMID 20628086.

The Fanconi anemia protein, FANCG, binds to the ERCC1-XPF endonuclease via its tetratricopeptide repeats and the central domain of ERCC1. Wang C, et al. Biochemistry, 2010 Jul 6. PMID 20518486.

Comprehensive screen of genetic variation in DNA repair pathway genes and postmenopausal breast cancer risk. Monsees GM, et al. Breast Cancer Res Treat, 2010 May 23. PMID 20496165.

Genetic variation in 3-hydroxy-3-methylglutaryl CoA reductase modifies the chemopreventive activity of statins for colorectal cancer. Lipkin SM, et al. Cancer Prev Res (Phila), 2010 May. PMID 20403997.

A histone-fold complex and FANCM form a conserved DNA-remodeling complex to maintain genome stability. Yan Z, et al. Mol Cell, 2010 Mar 26. PMID 20347428.