

Goat Anti-LASP1 (N Terminus) Antibody
Peptide-affinity purified goat antibody
Catalog # AF1613a**Specification**

Goat Anti-LASP1 (N Terminus) Antibody - Product Information

Application	WB
Primary Accession	Q14847
Other Accession	NP_006139 , 3927 , 16796 (mouse) , 29278 (rat)
Reactivity	Human
Predicted	Mouse, Rat, Pig, Dog
Host	Goat
Clonality	Polyclonal
Concentration	100ug/200ul
Isotype	IgG
Calculated MW	29717

Goat Anti-LASP1 (N Terminus) Antibody - Additional Information**Gene ID** 3927**Other Names**

LIM and SH3 domain protein 1, LASP-1, Metastatic lymph node gene 50 protein, MLN 50, LASP1, MLN50

Format

0.5 mg IgG/ml in Tris saline (20mM Tris pH7.3, 150mM NaCl), 0.02% sodium azide, with 0.5% bovine serum albumin

Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

Goat Anti-LASP1 (N Terminus) Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

Goat Anti-LASP1 (N Terminus) Antibody - Protein Information**Name** LASP1**Synonyms** MLN50**Function**

Plays an important role in the regulation of dynamic actin- based, cytoskeletal activities. Agonist-dependent changes in LASP1 phosphorylation may also serve to regulate actin-associated ion transport activities, not only in the parietal cell but also in certain other F-actin-rich secretory

epithelial cell types (By similarity).

Cellular Location

Cytoplasm, cell cortex. Cytoplasm, cytoskeleton. Note=Associated with the F-actin rich cortical cytoskeleton.

Goat Anti-LASP1 (N Terminus) Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Goat Anti-LASP1 (N Terminus) Antibody - Images



AF1613a staining (0.1 µg/ml) of Human Brain (Cerebellum) extracts (RIPA buffer, 35 µg total protein per lane). Primary incubated for 1 hour. Detected by western blot using chemiluminescence.

Goat Anti-LASP1 (N Terminus) Antibody - Background

This gene encodes a member of a LIM protein subfamily characterized by a LIM motif and a domain of Src homology region 3. The encoded protein functions as an actin-binding protein and possibly in cytoskeletal organization.

Goat Anti-LASP1 (N Terminus) Antibody - References

Nuclear localisation of LASP-1 correlates with poor long-term survival in female breast cancer. Frietsch JJ, et al. Br J Cancer, 2010 May 25. PMID 20461080.
Personalized smoking cessation: interactions between nicotine dose, dependence and quit-success genotype score. Rose JE, et al. Mol Med, 2010 Jul-Aug. PMID 20379614.
Integrin-dependent translocation of LASP-1 to the cytoskeleton of activated platelets correlates with LASP-1 phosphorylation at tyrosine 171 by Src-kinase. Traenka J, et al. Thromb Haemost, 2009 Sep. PMID 19718473.

Proteomic identification of LASP-1 down-regulation after RNAi urokinase silencing in human hepatocellular carcinoma cells. Salvi A, et al. Neoplasia, 2009 Feb. PMID 19177205.
LIM and SH3 protein 1 (Lasp1) is a novel p53 transcriptional target involved in hepatocellular carcinoma. Wang B, et al. J Hepatol, 2009 Mar. PMID 19155088.