

Goat Anti-Melanocortin 3 Receptor Antibody
Peptide-affinity purified goat antibody
Catalog # AF1666a**Specification**

Goat Anti-Melanocortin 3 Receptor Antibody - Product Information

Application	WB, E
Primary Accession	P41968
Other Accession	NP_063941 , 4159
Reactivity	Human
Host	Goat
Clonality	Polyclonal
Concentration	100ug/200ul
Isotype	IgG
Calculated MW	36043

Goat Anti-Melanocortin 3 Receptor Antibody - Additional Information**Gene ID** 4159**Other Names**

Melanocortin receptor 3, MC3-R, MC3R

Dilution

WB~~1:1000

E~~N/A

Format

0.5 mg IgG/ml in Tris saline (20mM Tris pH7.3, 150mM NaCl), 0.02% sodium azide, with 0.5% bovine serum albumin

Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

Goat Anti-Melanocortin 3 Receptor Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

Goat Anti-Melanocortin 3 Receptor Antibody - Protein Information**Name** MC3R**Function**

Receptor for MSH (alpha, beta and gamma) and ACTH. This receptor is mediated by G proteins which activate adenylate cyclase. Required for expression of anticipatory patterns of activity and wakefulness during periods of limited nutrient availability and for the normal regulation of

circadian clock activity in the brain.

Cellular Location

Cell membrane; Multi-pass membrane protein.

Tissue Location

Brain, placental, and gut tissues.


Goat Anti-Melanocortin 3 Receptor Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Goat Anti-Melanocortin 3 Receptor Antibody - Images

AF1666a (1 µg/ml) staining of Human Brain lysate (35 µg protein in RIPA buffer). Primary incubation was 1 hour. Detected by chemiluminescence.



250kDa
150kDa
100kDa
75kDa
50kDa
37kDa
25kDa
20kDa
15kDa
10kDa

EB06726 (1 μ g/ml) staining of Human Brain lysate (35 μ g protein in RIPA buffer). Detected by chemiluminescence.

Goat Anti-Melanocortin 3 Receptor Antibody - Background

This gene encodes a G-protein-coupled receptor for melanocyte-stimulating hormone and adrenocorticotrophic hormone that is expressed in tissues other than the adrenal cortex and melanocytes. This gene maps to the same region as the locus for benign neonatal epilepsy. Mice deficient for this gene have increased fat mass despite decreased food intake, suggesting a role for this gene product in the regulation of energy homeostasis. Mutations in this gene are associated with a susceptibility to obesity in humans.

Goat Anti-Melanocortin 3 Receptor Antibody - References

Physiogenomic analysis of statin-treated patients: domain-specific counter effects within the ACACB gene on low-density lipoprotein cholesterol? Ruaño G, et al. Pharmacogenomics, 2010 Jul. PMID 20602615.
Identification of Three Novel Genetic Variants in the Melanocortin-3 Receptor of Obese Children. Zegers D, et al. Obesity (Silver Spring), 2010 Jun 10. PMID 20539302.
Association study of 182 candidate genes in anorexia nervosa. Pinheiro AP, et al. Am J Med Genet B Neuropsychiatr Genet, 2010 Jul. PMID 20468064.
Melanocortin-3 receptor gene variants: association with childhood obesity and eating behavior in Chilean families. Obregón AM, et al. Nutrition, 2010 Jul-Aug. PMID 20144537.
Novel binding motif of ACTH analogues at the melanocortin receptors. Yang Y, et al. Biochemistry, 2009 Oct 20. PMID 19743876.