

### Goat Anti-MSX1 Antibody

Peptide-affinity purified goat antibody Catalog # AF1687a

### Specification

# **Goat Anti-MSX1 Antibody - Product Information**

Application Primary Accession Other Accession Reactivity Predicted Host Clonality Concentration Isotype Calculated MW

WB, IHC, E <u>P28360</u> <u>NP\_002439</u>, <u>4487</u>, <u>17701 (mouse)</u> Human Mouse, Rat, Pig Goat Polyclonal 100ug/200ul IgG 31496

# **Goat Anti-MSX1 Antibody - Additional Information**

Gene ID 4487

**Other Names** Homeobox protein MSX-1, Homeobox protein Hox-7, Msh homeobox 1-like protein, MSX1, HOX7

**Dilution** WB~~1:1000 IHC~~1:100~500 E~~N/A

Format

0.5 mg lgG/ml in Tris saline (20mM Tris pH7.3, 150mM NaCl), 0.02% sodium azide, with 0.5% bovine serum albumin

Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

**Precautions** Goat Anti-MSX1 Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

# **Goat Anti-MSX1 Antibody - Protein Information**

Name MSX1 (<u>HGNC:7391</u>)

Function

Acts as a transcriptional repressor (By similarity). Capable of transcription autoinactivation (By



similarity). Binds to the consensus sequence 5'-C/GTAAT-3' in downstream activin regulatory elements (DARE) in the gene promoter, thereby repressing the transcription of CGA/alpha-GSU and GNRHR (By similarity). Represses transcription of myoblast differentiation factors (By similarity). Binds to core enhancer regions in target gene promoters of myoblast differentiation factors with binding specificity facilitated by interaction with PIAS1 (By similarity). Regulates, in a stage-specific manner, a developmental program of gene expression in the fetal tooth bud that controls odontoblast differentiation and proliferation of dental mesenchymal cells (By similarity). At the bud stage, required for mesenchymal molar tooth bud development via facilitating reciprocal signaling between dental epithelial and mesenchymal cells (By similarity). May also regulate expression of Wnt antagonists such as DKK2 and SFPR2 in the developing tooth mesenchyme (By similarity). Required for BMP4 expression in dental mesenchyme cells (By similarity). Also, in response to BMP4, required for BMP4 expression in neighboring dental epithelial cells (By similarity). Required for maximal FGF4-induced expression of SDC1 in dental mesenchyme cells (By similarity). Also in response to SDC1, required for SDC1 expression in neighboring dental epithelial cells (By similarity). At the early bell stage, acts to drive proliferation of dental mesenchyme cells, however during the late bell stage acts as an homeostatic regulator of the cell cycle (By similarity). Regulates proliferation and inhibits premature mesenchymal odontogenesis during the bell stage via inhibition of the Wnt signaling component CTNNB1 and subsequent repression of the odontoblast differentiation factors BMP2, BMP4, LEF1, ALPL and BGLAP/OCN (By similarity). Additionally, required for correct development and fusion of the palatal shelves and embryonic mandibular formation (By similarity). Plays a role in embryonic bone formation of the middle ear, skull and nasal bones (By similarity). Required for correct formation and thickness of the nail plate (By similarity). May play a role in limb-pattern formation (By similarity).

#### **Cellular Location**

Nucleus {ECO:0000250|UniProtKB:P13297}. Note=Interaction with PIAS1 is required for localization to the nuclear periphery (By similarity). {ECO:0000250|UniProtKB:P13297}

#### Goat Anti-MSX1 Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- <u>Western Blot</u>
- Blocking Peptides
- <u>Dot Blot</u>
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- <u>Cell Culture</u>

Goat Anti-MSX1 Antibody - Images

	250kDa 150kDa 100kDa 75kDa	
	50kDa	
	37kDa	
-		
	25kDa	
	20kDa	
	15kDa	

AF1687a (0.3  $\mu$ g/ml) staining of Human Prostate lysate (35  $\mu$ g protein in RIPA buffer). Primary incubation was 1 hour. Detected by chemiluminescence.



AF1687a (2.5  $\mu$ g/ml) staining of paraffin embedded Human Prostate. Steamed antigen retrieval with citrate buffer pH 6, AP-staining.

# Goat Anti-MSX1 Antibody - Background

This gene encodes a member of the muscle segment homeobox gene family. The encoded protein functions as a transcriptional repressor during embryogenesis through interactions with components of the core transcription complex and other homeoproteins. It may also have roles in limb-pattern formation, craniofacial development, particularly odontogenesis, and tumor growth inhibition. Mutations in this gene, which was once known as homeobox 7, have been associated with nonsyndromic cleft lip with or without cleft palate 5, Witkop syndrome, Wolf-Hirschom syndrome, and autosomoal dominant hypodontia.

#### Goat Anti-MSX1 Antibody - References

Genetic variants in COL2A1, COL11A2, and IRF6 contribute risk to nonsyndromic cleft palate. Nikopensius T, et al. Birth Defects Res A Clin Mol Teratol, 2010 Jul 29. PMID 20672350. The c.469+46\_56del mutation in the homeobox MSX1 gene-A novel risk factor in breast cancer? Sliwinski T, et al. Cancer Epidemiol, 2010 Jul 16. PMID 20638926.

Parent-of-origin effects for MSX1 in a Chilean population with nonsyndromic cleft lip/palate. Suazo J, et al. Am J Med Genet A, 2010 Aug. PMID 20635363.

Maternal genes and facial clefts in offspring: a comprehensive search for genetic associations in two population-based cleft studies from Scandinavia. Jugessur A, et al. PLoS One, 2010 Jul 9. PMID 20634891.



MTHFR and MSX1 contribute to the risk of nonsyndromic cleft lip/palate. Jagom gi T, et al. Eur J Oral Sci, 2010 Jun. PMID 20572854.