

Goat Anti-MURF3 / TRIM54 Antibody

Peptide-affinity purified goat antibody Catalog # AF1698a

Specification

Goat Anti-MURF3 / TRIM54 Antibody - Product Information

Application WB
Primary Accession O9BYV2

Other Accession NP 912730, 57159

Reactivity
Predicted
Dog
Host
Clonality
Human
Dog
Goat
Polyclonal

Concentration 100ug/200ul Isotype IgG
Calculated MW 40301

Goat Anti-MURF3 / TRIM54 Antibody - Additional Information

Gene ID 57159

Other Names

Tripartite motif-containing protein 54, Muscle-specific RING finger protein, MuRF, Muscle-specific RING finger protein 3, MuRF-3, MuRF3, RING finger protein 30, TRIM54, MURF, MURF3, RNF30

Format

0.5~mg lgG/ml in Tris saline (20mM Tris pH7.3, 150mM NaCl), 0.02% sodium azide, with 0.5% bovine serum albumin

Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

Goat Anti-MURF3 / TRIM54 Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

Goat Anti-MURF3 / TRIM54 Antibody - Protein Information

Name TRIM54

Synonyms MURF, MURF3, RNF30

Function

May bind and stabilize microtubules during myotubes formation.

Cellular Location



Cytoplasm, cytoskeleton. Cytoplasm, myofibril, sarcomere, Z line. Note=Associates with microtubules. Localizes to the Z-lines in skeletal muscles (By similarity).

Tissue Location


Specifically expressed in heart and skeletal muscle.

Goat Anti-MURF3 / TRIM54 Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- <u>Immunohistochemistry</u>
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

Goat Anti-MURF3 / TRIM54 Antibody - Images



AF1698a staining (1ug/ml) of human muscle lysate (RIPA buffer, 35ug totalprotein per lane). Primary incubated for 1 hour. Detected by western blotusing chemiluminescence.

Goat Anti-MURF3 / TRIM54 Antibody - Background

The protein encoded by this gene contains a RING finger motif and is highly similar to the ring finger proteins RNF28/MURF1 and RNF29/MURF2. In vitro studies demonstrated that this protein, RNF28, and RNF29 form heterodimers, which may be important for the regulation of titin kinase and microtubule-dependent signal pathways in striated muscles. Alternatively spliced transcript variants encoding distinct isoforms have been reported.

Goat Anti-MURF3 / TRIM54 Antibody - References

Diversification of transcriptional modulation: large-scale identification and characterization of putative alternative promoters of human genes. Kimura K, et al. Genome Res, 2006 Jan. PMID 16344560.

Generation and annotation of the DNA sequences of human chromosomes 2 and 4. Hillier LW, et al. Nature, 2005 Apr 7. PMID 15815621.

Generation and initial analysis of more than 15,000 full-length human and mouse cDNA sequences.





Tel: 858.875.1900 Fax: 858.875.1999

Strausberg RL, et al. Proc Natl Acad Sci U S A, 2002 Dec 24. PMID 12477932. Muscle-specific RING finger-1 interacts with titin to regulate sarcomeric M-line and thick filament structure and may have nuclear functions via its interaction with glucocorticoid modulatory element binding protein-1. McElhinny AS, et al. J Cell Biol, 2002 Apr 1. PMID 11927605. Identification of muscle specific ring finger proteins as potential regulators of the titin kinase domain. Centner T, et al. J Mol Biol, 2001 Mar 2. PMID 11243782.