

Goat Anti-OGT Antibody
Peptide-affinity purified goat antibody
Catalog # AF1752a

Specification

Goat Anti-OGT Antibody - Product Information

Application	WB, IHC, E
Primary Accession	O15294
Other Accession	NP_858059 , 8473 , 108155 (mouse) , 26295 (rat)
Reactivity	Mouse, Rat
Predicted	Human, Dog
Host	Goat
Clonality	Polyclonal
Concentration	100ug/200ul
Isotype	IgG
Calculated MW	116925

Goat Anti-OGT Antibody - Additional Information

Gene ID 8473

Other Names

UDP-N-acetylglucosamine--peptide N-acetylglucosaminyltransferase 110 kDa subunit, 2.4.1.255, O-GlcNAc transferase subunit p110, O-linked N-acetylglucosamine transferase 110 kDa subunit, OGT, OGT

Dilution

WB~~1:1000
IHC~~1:100~500
E~~N/A

Format

0.5 mg IgG/ml in Tris saline (20mM Tris pH7.3, 150mM NaCl), 0.02% sodium azide, with 0.5% bovine serum albumin

Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

Goat Anti-OGT Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

Goat Anti-OGT Antibody - Protein Information

Name OGT {ECO:0000303|PubMed:11773972, ECO:0000312|HGNC:HGNC:8127}

Function

Catalyzes the transfer of a single N-acetylglucosamine from UDP-GlcNAc to a serine or threonine residue in cytoplasmic and nuclear proteins resulting in their modification with a beta-linked N-acetylglucosamine (O-GlcNAc) (PubMed: [12150998](http://www.uniprot.org/citations/12150998), PubMed: [15361863](http://www.uniprot.org/citations/15361863), PubMed: [19451179](http://www.uniprot.org/citations/19451179), PubMed: [20018868](http://www.uniprot.org/citations/20018868), PubMed: [21240259](http://www.uniprot.org/citations/21240259), PubMed: [21285374](http://www.uniprot.org/citations/21285374), PubMed: [23103939](http://www.uniprot.org/citations/23103939), PubMed: [26237509](http://www.uniprot.org/citations/26237509), PubMed: [26369908](http://www.uniprot.org/citations/26369908), PubMed: [26678539](http://www.uniprot.org/citations/26678539), PubMed: [27713473](http://www.uniprot.org/citations/27713473), PubMed: [37541260](http://www.uniprot.org/citations/37541260), PubMed: [37962578](http://www.uniprot.org/citations/37962578)). Glycosylates a large and diverse number of proteins including histone H2B, AKT1, AMPK, ATG4B, CAPRIN1, EZH2, FNIP1, GSDMD, KRT7, LMNA, LMNB1, LMNB2, RPTOR, HOXA1, PFKL, KMT2E/MLL5, MAPT/TAU, TET2, RBL2, RET, NOD2 and HCFC1 (PubMed: [19451179](http://www.uniprot.org/citations/19451179), PubMed: [20200153](http://www.uniprot.org/citations/20200153), PubMed: [21285374](http://www.uniprot.org/citations/21285374), PubMed: [22923583](http://www.uniprot.org/citations/22923583), PubMed: [23353889](http://www.uniprot.org/citations/23353889), PubMed: [24474760](http://www.uniprot.org/citations/24474760), PubMed: [26237509](http://www.uniprot.org/citations/26237509), PubMed: [26369908](http://www.uniprot.org/citations/26369908), PubMed: [26678539](http://www.uniprot.org/citations/26678539), PubMed: [27527864](http://www.uniprot.org/citations/27527864), PubMed: [30699359](http://www.uniprot.org/citations/30699359), PubMed: [34074792](http://www.uniprot.org/citations/34074792), PubMed: [34667079](http://www.uniprot.org/citations/34667079), PubMed: [37541260](http://www.uniprot.org/citations/37541260), PubMed: [37962578](http://www.uniprot.org/citations/37962578)). Can regulate their cellular processes via cross-talk between glycosylation and phosphorylation or by affecting proteolytic processing (PubMed: [21285374](http://www.uniprot.org/citations/21285374)). Involved in insulin resistance in muscle and adipocyte cells via glycosylating insulin signaling components and inhibiting the 'Thr-308' phosphorylation of AKT1, enhancing IRS1 phosphorylation and attenuating insulin signaling (By similarity). Involved in glycolysis regulation by mediating glycosylation of 6-phosphofructokinase PFKL, inhibiting its activity (PubMed: [22923583](http://www.uniprot.org/citations/22923583)). Plays a key role in chromatin structure by mediating O-GlcNAcylation of 'Ser-112' of histone H2B: recruited to CpG-rich transcription start sites of active genes via its interaction with TET proteins (TET1, TET2 or TET3) (PubMed: [22121020](http://www.uniprot.org/citations/22121020), PubMed: [23353889](http://www.uniprot.org/citations/23353889)). As part of the NSL complex indirectly involved in acetylation of nucleosomal histone H4 on several lysine residues (PubMed: [20018852](http://www.uniprot.org/citations/20018852)). O-GlcNAcylation of 'Ser-75' of EZH2 increases its stability, and facilitating the formation of H3K27me3 by the PRC2/EED-EZH2 complex (PubMed: [24474760](http://www.uniprot.org/citations/24474760)). Stabilizes KMT2E/MLL5 by mediating its glycosylation, thereby preventing KMT2E/MLL5 ubiquitination (PubMed: [26678539](http://www.uniprot.org/citations/26678539)). Regulates circadian oscillation of the clock genes and glucose homeostasis in the liver (By similarity). Stabilizes clock proteins BMAL1 and CLOCK through O-glycosylation, which prevents their ubiquitination and subsequent degradation (By similarity). Promotes the CLOCK-BMAL1-mediated transcription of genes in the negative loop of the circadian clock such as

PER1/2 and CRY1/2. O-glycosylates HCFC1 and regulates its proteolytic processing and transcriptional activity (PubMed:21285374, PubMed:28302723, PubMed:28584052). Component of a THAP1/THAP3-HCFC1-OGT complex that is required for the regulation of the transcriptional activity of RRM1 (PubMed:20200153). Regulates mitochondrial motility in neurons by mediating glycosylation of TRAK1 (By similarity). Promotes autophagy by mediating O-glycosylation of ATG4B (PubMed:27527864). Acts as a regulator of mTORC1 signaling by mediating O-glycosylation of RPTOR and FNIP1: O-GlcNAcylation of RPTOR in response to glucose sufficiency promotes activation of the mTORC1 complex (PubMed:30699359, PubMed:37541260).

Cellular Location

Nucleus. Cytoplasm. Note=Predominantly localizes to the nucleus (PubMed:26678539). Translocates into the nucleus via association with importin KPNA1 (PubMed:27713473) [Isoform 3]: Cytoplasm. Nucleus. Cell membrane {ECO:0000250|UniProtKB:P56558}. Mitochondrion membrane {ECO:0000250|UniProtKB:P56558}. Cell projection {ECO:0000250|UniProtKB:P56558}. Note=Mostly in the nucleus. Retained in the nucleus via interaction with HCFC1 (PubMed:21285374). After insulin induction, translocated from the nucleus to the cell membrane via phosphatidylinositide binding. Colocalizes with AKT1 at the plasma membrane. TRAK1 recruits this protein to mitochondria. In the absence of TRAK1, localizes in cytosol and nucleus (By similarity) {ECO:0000250|UniProtKB:P56558, ECO:0000269|PubMed:21285374}

Tissue Location


Highly expressed in pancreas and to a lesser extent in skeletal muscle, heart, brain and placenta. Present in trace amounts in lung and liver.


Goat Anti-OGT Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.


- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Goat Anti-OGT Antibody - Images





AF1752a (0.05 $\mu\text{g/ml}$) staining of Rat Pancreas lysate (35 μg protein in RIPA buffer). Primary incubation was 1 hour. Detected by chemiluminescence.



AF1752a (5 $\mu\text{g/ml}$) staining of paraffin embedded Human Cortex. Steamed antigen retrieval with citrate buffer pH 6, AP-staining.