

**Goat Anti-ribosomal protein L8 Antibody**  
**Peptide-affinity purified goat antibody**  
**Catalog # AF1928a****Specification**

---

**Goat Anti-ribosomal protein L8 Antibody - Product Information**

Application	WB, IHC, E
Primary Accession	<a href="#">P62917</a>
Other Accession	<a href="#">NP_150644</a> , <a href="#">6132</a> , <a href="#">26961 (mouse)</a> , <a href="#">26962 (rat)</a>
Reactivity	Human
Predicted	Mouse, Rat, Dog
Host	Goat
Clonality	Polyclonal
Concentration	100ug/200ul
Isotype	IgG
Calculated MW	28025

**Goat Anti-ribosomal protein L8 Antibody - Additional Information****Gene ID** 6132**Other Names**

60S ribosomal protein L8, RPL8

**Dilution**

WB~~1:1000

IHC~~1:100~500

E~~N/A

**Format**

0.5 mg IgG/ml in Tris saline (20mM Tris pH7.3, 150mM NaCl), 0.02% sodium azide, with 0.5% bovine serum albumin

**Storage**

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

**Precautions**

Goat Anti-ribosomal protein L8 Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

**Goat Anti-ribosomal protein L8 Antibody - Protein Information****Name** RPL8**Function**

Component of the large ribosomal subunit. The ribosome is a large ribonucleoprotein complex

responsible for the synthesis of proteins in the cell.

#### **Cellular Location**

Cytoplasm.

#### **Goat Anti-ribosomal protein L8 Antibody - Protocols**


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

#### **Goat Anti-ribosomal protein L8 Antibody - Images**



AF1928a (0.01 µg/ml) staining of HepG2 cell lysate (35 µg protein in RIPA buffer). Primary incubation was 1 hour. Detected by chemiluminescence.



AF1928a (2.5 µg/ml) staining of paraffin embedded Human Cerebral Cortex. Steamed antigen retrieval with citrate buffer pH 6, AP-staining.

#### **Goat Anti-ribosomal protein L8 Antibody - Background**

Ribosomes, the organelles that catalyze protein synthesis, consist of a small 40S subunit and a large 60S subunit. Together these subunits are composed of 4 RNA species and approximately 80

structurally distinct proteins. This gene encodes a ribosomal protein that is a component of the 60S subunit. The protein belongs to the L2P family of ribosomal proteins. It is located in the cytoplasm. In rat, the protein associates with the 5.8S rRNA, very likely participates in the binding of aminoacyl-tRNA, and is a constituent of the elongation factor 2-binding site at the ribosomal subunit interface. Alternatively spliced transcript variants encoding the same protein exist. As is typical for genes encoding ribosomal proteins, there are multiple processed pseudogenes of this gene dispersed through the genome.

#### **Goat Anti-ribosomal protein L8 Antibody - References**

Molecular characterization of the response to chemotherapy in conventional osteosarcomas: predictive value of HSD17B10 and IFITM2. Salas S, et al. *Int J Cancer*, 2009 Aug 15. PMID 19449377.

A human protein-protein interaction network: a resource for annotating the proteome. Stelzl U, et al. *Cell*, 2005 Sep 23. PMID 16169070.

Nucleolar proteome dynamics. Andersen JS, et al. *Nature*, 2005 Jan 6. PMID 15635413.

Immunoaffinity profiling of tyrosine phosphorylation in cancer cells. Rush J, et al. *Nat Biotechnol*, 2005 Jan. PMID 15592455.

The status, quality, and expansion of the NIH full-length cDNA project: the Mammalian Gene Collection (MGC). Gerhard DS, et al. *Genome Res*, 2004 Oct. PMID 15489334.