

**Goat Anti-STAM2 Antibody**  
**Peptide-affinity purified goat antibody**  
**Catalog # AF2038a****Specification**

---

**Goat Anti-STAM2 Antibody - Product Information**

Application	WB, E
Primary Accession	<a href="#">O75886</a>
Other Accession	<a href="#">NP_005834</a> , <a href="#">10254</a> , <a href="#">56324 (mouse)</a>
Reactivity	Human
Predicted	Mouse, Dog
Host	Goat
Clonality	Polyclonal
Concentration	100ug/200ul
Isotype	IgG
Calculated MW	58164

**Goat Anti-STAM2 Antibody - Additional Information****Gene ID** 10254**Other Names**

Signal transducing adapter molecule 2, STAM-2, Hrs-binding protein, STAM2, HBP

**Dilution**

WB~~1:1000

E~~N/A

**Format**

0.5 mg IgG/ml in Tris saline (20mM Tris pH7.3, 150mM NaCl), 0.02% sodium azide, with 0.5% bovine serum albumin

**Storage**

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

**Precautions**

Goat Anti-STAM2 Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

**Goat Anti-STAM2 Antibody - Protein Information****Name** STAM2**Synonyms** HBP**Function**

Involved in intracellular signal transduction mediated by cytokines and growth factors. Upon IL-2 and GM-CSF stimulation, it plays a role in signaling leading to DNA synthesis and MYC induction. May also play a role in T-cell development. Involved in down-regulation of receptor tyrosine kinase via multivesicular body (MVBs) when complexed with HGS (ESCRT-0 complex). The ESCRT-0 complex binds ubiquitin and acts as a sorting machinery that recognizes ubiquitinated receptors and transfers them to further sequential lysosomal sorting/trafficking processes (By similarity).

**Cellular Location**

Cytoplasm. Early endosome membrane; Peripheral membrane protein; Cytoplasmic side

**Tissue Location**

Ubiquitously expressed.

**Goat Anti-STAM2 Antibody - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

**Goat Anti-STAM2 Antibody - Images**

AF2038a (1 µg/ml) staining of Human Spleen lysate (35 µg protein in RIPA buffer). Primary incubation was 1 hour. Detected by chemiluminescence.

**Goat Anti-STAM2 Antibody - Background**

The protein encoded by this gene is closely related to STAM, an adaptor protein involved in the downstream signaling of cytokine receptors, both of which contain a SH3 domain and the immunoreceptor tyrosine-based activation motif (ITAM). Similar to STAM, this protein acts downstream of JAK kinases, and is phosphorylated in response to cytokine stimulation. This protein and STAM thus are thought to exhibit compensatory effects on the signaling pathway downstream of JAK kinases upon cytokine stimulation.

## **Goat Anti-STAM2 Antibody - References**

PTP1B targets the endosomal sorting machinery: dephosphorylation of regulatory sites on the endosomal sorting complex required for transport component STAM2. Stuible M, et al. J Biol Chem, 2010 Jul 30. PMID 20504764.

An empirical framework for binary interactome mapping. Venkatesan K, et al. Nat Methods, 2009 Jan. PMID 19060904.

STAM adaptor proteins interact with COPII complexes and function in ER-to-Golgi trafficking. Rismanchi N, et al. Traffic, 2009 Feb. PMID 19054391.

E3-independent monoubiquitination of ubiquitin-binding proteins. Hoeller D, et al. Mol Cell, 2007 Jun 22. PMID 17588522.

Rin1 interacts with signal-transducing adaptor molecule (STAM) and mediates epidermal growth factor receptor trafficking and degradation. Kong C, et al. J Biol Chem, 2007 May 18. PMID 17403676.