

**Goat Anti-STAT3 (isoform 1 and 2) Antibody**  
**Peptide-affinity purified goat antibody**  
**Catalog # AF2040a****Specification**

---

**Goat Anti-STAT3 (isoform 1 and 2) Antibody - Product Information**

Application	IHC
Primary Accession	<a href="#">P40763</a>
Other Accession	<a href="#">NP_644805</a> , <a href="#">6774</a> , <a href="#">20848 (mouse)</a> , <a href="#">25125 (rat)</a>
Reactivity	Human, Mouse, Rat
Predicted	Pig, Dog, Cow
Host	Goat
Clonality	Polyclonal
Concentration	100ug/200ul
Isotype	IgG
Calculated MW	88068

**Goat Anti-STAT3 (isoform 1 and 2) Antibody - Additional Information****Gene ID** 6774**Other Names**

Signal transducer and activator of transcription 3, Acute-phase response factor, STAT3, APRF

**Format**

0.5 mg IgG/ml in Tris saline (20mM Tris pH7.3, 150mM NaCl), 0.02% sodium azide, with 0.5% bovine serum albumin

**Storage**

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

**Precautions**

Goat Anti-STAT3 (isoform 1 and 2) Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

**Goat Anti-STAT3 (isoform 1 and 2) Antibody - Protein Information****Name** STAT3 {ECO:0000303|PubMed:9630560, ECO:0000312|HGNC:HGNC:11364}**Function**

Signal transducer and transcription activator that mediates cellular responses to interleukins, KITLG/SCF, LEP and other growth factors (PubMed:&lt;a href="http://www.uniprot.org/citations/10688651" target="\_blank"&gt;10688651&lt;/a&gt;, PubMed:&lt;a href="http://www.uniprot.org/citations/12359225" target="\_blank"&gt;12359225&lt;/a&gt;, PubMed:&lt;a href="http://www.uniprot.org/citations/12873986" target="\_blank"&gt;12873986&lt;/a&gt;, PubMed:&lt;a href="http://www.uniprot.org/citations/15194700" target="\_blank"&gt;15194700&lt;/a&gt;, PubMed:&lt;a

[16285960](http://www.uniprot.org/citations/16285960), PubMed: [15653507](http://www.uniprot.org/citations/15653507), PubMed: [17344214](http://www.uniprot.org/citations/17344214), PubMed: [18242580](http://www.uniprot.org/citations/18242580), PubMed: [18782771](http://www.uniprot.org/citations/18782771), PubMed: [22306293](http://www.uniprot.org/citations/22306293), PubMed: [23084476](http://www.uniprot.org/citations/23084476), PubMed: [32929201](http://www.uniprot.org/citations/32929201), PubMed: [28262505](http://www.uniprot.org/citations/28262505)). Once activated, recruits coactivators, such as NCOA1 or MED1, to the promoter region of the target gene (PubMed: [16285960](http://www.uniprot.org/citations/16285960), PubMed: [15653507](http://www.uniprot.org/citations/15653507), PubMed: [17344214](http://www.uniprot.org/citations/17344214), PubMed: [18782771](http://www.uniprot.org/citations/18782771), PubMed: [28262505](http://www.uniprot.org/citations/28262505), PubMed: [32929201](http://www.uniprot.org/citations/32929201)). May mediate cellular responses to activated FGFR1, FGFR2, FGFR3 and FGFR4 (PubMed: [12873986](http://www.uniprot.org/citations/12873986)). Upon activation of IL6ST/gp130 signaling by interleukin-6 (IL6), binds to the IL6-responsive elements identified in the promoters of various acute-phase protein genes (PubMed: [12359225](http://www.uniprot.org/citations/12359225)). Activated by IL31 through IL31RA (PubMed: [15194700](http://www.uniprot.org/citations/15194700)). Acts as a regulator of inflammatory response by regulating differentiation of naive CD4(+) T-cells into T-helper Th17 or regulatory T-cells (Treg): acetylation promotes its transcription activity and cell differentiation while deacetylation and oxidation of lysine residues by LOXL3 inhibits differentiation (PubMed: [28262505](http://www.uniprot.org/citations/28262505), PubMed: [28065600](http://www.uniprot.org/citations/28065600)). Involved in cell cycle regulation by inducing the expression of key genes for the progression from G1 to S phase, such as CCND1 (PubMed: [17344214](http://www.uniprot.org/citations/17344214)). Mediates the effects of LEP on melanocortin production, body energy homeostasis and lactation (By similarity). May play an apoptotic role by transactivating BIRC5 expression under LEP activation (PubMed: [18242580](http://www.uniprot.org/citations/18242580)). Cytoplasmic STAT3 represses macroautophagy by inhibiting EIF2AK2/PKR activity (PubMed: [23084476](http://www.uniprot.org/citations/23084476)). Plays a crucial role in basal beta cell functions, such as regulation of insulin secretion (By similarity). Following JAK/STAT signaling activation and as part of a complex with NFATC3 and NFATC4, binds to the alpha-beta E4 promoter region of CRYAB and activates transcription in cardiomyocytes (By similarity).

### Cellular Location

Cytoplasm. Nucleus. Note=Shuttles between the nucleus and the cytoplasm. Translocated into the nucleus upon tyrosine phosphorylation and dimerization, in response to signaling by activated FGFR1, FGFR2, FGFR3 or FGFR4 (PubMed:16285960, PubMed:15653507). Constitutive nuclear presence is independent of tyrosine phosphorylation. Predominantly present in the cytoplasm without stimuli. Upon leukemia inhibitory factor (LIF) stimulation, accumulates in the nucleus. The complex composed of BART and ARL2 plays an important role in the nuclear translocation and retention of STAT3. Identified in a complex with LYN and PAG1. Translocates to the nucleus in the presence of EDN1 (By similarity). {ECO:0000250|UniProtKB:P52631, ECO:0000269|PubMed:15653507, ECO:0000269|PubMed:16285960}

### Tissue Location


Heart, brain, placenta, lung, liver, skeletal muscle, kidney and pancreas. Expressed in naive CD4(+) T cells as well as T-helper Th17, Th1 and Th2 cells (PubMed:31899195)

## Goat Anti-STAT3 (isoform 1 and 2) Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

## Goat Anti-STAT3 (isoform 1 and 2) Antibody - Images



AF2040a (4 µg/ml) staining of paraffin embedded Mouse Thymus. Steamed antigen retrieval with Tris/EDTA buffer pH 9, HRP-staining

## Goat Anti-STAT3 (isoform 1 and 2) Antibody - Background

The protein encoded by this gene is a member of the STAT protein family. In response to cytokines and growth factors, STAT family members are phosphorylated by the receptor associated kinases, and then form homo- or heterodimers that translocate to the cell nucleus where they act as transcription activators. This protein is activated through phosphorylation in response to various cytokines and growth factors including IFNs, EGF, IL5, IL6, HGF, LIF and BMP2. This protein mediates the expression of a variety of genes in response to cell stimuli, and thus plays a key role in many cellular processes such as cell growth and apoptosis. The small GTPase Rac1 has been shown to bind and regulate the activity of this protein. PIAS3 protein is a specific inhibitor of this protein. Three alternatively spliced transcript variants encoding distinct isoforms have been described.

## Goat Anti-STAT3 (isoform 1 and 2) Antibody - References

Diagnostic approach to the hyper-IgE syndromes: immunologic and clinical key findings to differentiate hyper-IgE syndromes from atopic dermatitis. Schimke LF, et al. J Allergy Clin Immunol, 2010 Sep. PMID 20816194. Maternal genes and facial clefts in offspring: a comprehensive search for genetic associations in two population-based cleft studies from Scandinavia. Jugessur A, et al. PLoS One, 2010 Jul 9. PMID 20634891. Evaluation of candidate stromal epithelial cross-talk genes identifies association between risk of serous ovarian cancer and TERT, a cancer susceptibility hot-spot. Johnatty SE, et al. PLoS Genet, 2010 Jul 8. PMID 20628624. Variation at the NFATC2 Locus Increases the Risk of Thiazolinedione-Induced Edema in the Diabetes REDuction Assessment with ramipril and rosiglitazone Medication (DREAM) Study. Bailey SD, et al. Diabetes Care, 2010 Jul 13. PMID 20628086. Analysis of JAK2 and STAT3 polymorphisms in patients with ankylosing

spondylitis in Chinese Han population. Chen C, et al. Clin Immunol, 2010 Sep. PMID 20627814.