

**Goat Anti-U2AF1L4 Antibody**  
**Peptide-affinity purified goat antibody**  
**Catalog # AF2128a****Specification**

---

**Goat Anti-U2AF1L4 Antibody - Product Information**

|                   |  |
|-------------------|--|
| Application       | WB   |
| Primary Accession | <a href="#">Q8WU68</a>   |
| Other Accession   | <a href="#">NP_659424</a> , <a href="#">199746</a> , <a href="#">233073 (mouse)</a> , <a href="#">361542 (rat)</a> |
| Reactivity        | Human  |
| Predicted         | Mouse, Rat, Dog  |
| Host              | Goat   |
| Clonality         | Polyclonal   |
| Concentration     | 100ug/200ul  |
| Isotype           | IgG  |
| Calculated MW     | 25744  |

**Goat Anti-U2AF1L4 Antibody - Additional Information****Gene ID** 199746**Other Names**

Splicing factor U2AF 26 kDa subunit, U2 auxiliary factor 26, U2 small nuclear RNA auxiliary factor 1-like protein 4, U2AF1-like 4, U2(RNU2) small nuclear RNA auxiliary factor 1-like protein 3, U2 small nuclear RNA auxiliary factor 1-like protein 3, U2AF1-like protein 3, U2AF1L4, U2AF1-RS3, U2AF1L3

**Format**

0.5 mg IgG/ml in Tris saline (20mM Tris pH7.3, 150mM NaCl), 0.02% sodium azide, with 0.5% bovine serum albumin

**Storage**

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

**Precautions**

Goat Anti-U2AF1L4 Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

**Goat Anti-U2AF1L4 Antibody - Protein Information****Name** U2AF1L4**Synonyms** U2AF1-RS3, U2AF1L3**Function**

RNA-binding protein that function as a pre-mRNA splicing factor. Plays a critical role in both constitutive and enhancer- dependent splicing by mediating protein-protein interactions and protein-RNA interactions required for accurate 3'-splice site selection. Acts by enhancing the binding of U2AF2 to weak pyrimidine tracts. Also participates in the regulation of alternative pre-mRNA splicing. Activates exon 5 skipping of PTPRC during T-cell activation; an event reversed by GFI1. Binds to RNA at the AG dinucleotide at the 3'-splice site (By similarity). Shows a preference for AGC or AGA (By similarity).

#### Cellular Location

Nucleus {ECO:0000250|UniProtKB:Q8BGJ9}. Nucleus speckle {ECO:0000250|UniProtKB:Q8BGJ9}. Cytoplasm {ECO:0000250|UniProtKB:Q8BGJ9}. Note=Interaction with C1QBP is required for the nuclear translocation. Displays active nucleo-cytoplasmic shuttling. {ECO:0000250|UniProtKB:Q8BGJ9}

#### Tissue Location

Isoform 2 is widely expressed. Isoform 3 is highly expressed in heart, brain and lung, lower expressed in thymus and much lower expressed in peripheral blood leukocytes

### Goat Anti-U2AF1L4 Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

### Goat Anti-U2AF1L4 Antibody - Images



AF2128a (0.1 µg/ml) staining of Human Cerebellum lysate (35 µg protein in RIPA buffer). Primary incubation was 1 hour. Detected by chemiluminescence.

### Goat Anti-U2AF1L4 Antibody - References

Differential isoform expression and interaction with the P32 regulatory protein controls the subcellular localization of the splicing factor U2AF26. Heyd F, et al. J Biol Chem, 2008 Jul 11. PMID

18460468.

Cloning and characterization of a novel splice variant of human Rab18 gene (RAB18). Dou T, et al. DNA Seq, 2005 Jun. PMID 16147880.

Complete sequencing and characterization of 21,243 full-length human cDNAs. Ota T, et al. Nat Genet, 2004 Jan. PMID 14702039.

Generation and initial analysis of more than 15,000 full-length human and mouse cDNA sequences. Strausberg RL, et al. Proc Natl Acad Sci U S A, 2002 Dec 24. PMID 12477932.

Characterization of U2AF(6), a splicing factor related to U2AF(35). Shepard J, et al. Mol Cell Biol, 2002 Jan. PMID 11739736.