

Goat Anti-IL17ra (mouse) Antibody
Peptide-affinity purified goat antibody
Catalog # AF2172a**Specification**

Goat Anti-IL17ra (mouse) Antibody - Product Information

Application	WB
Primary Accession	Q60943.1
Other Accession	NP_032385 , 16172 (mouse) , 312679 (rat)
Reactivity	Rat
Predicted	Mouse
Host	Goat
Clonality	Polyclonal
Concentration	100ug/200ul
Isotype	IgG

Goat Anti-IL17ra (mouse) Antibody - Additional Information**Other Names**

interleukin 17 receptor A antibody, AW538159 antibody, Cdw217 antibody, Il17r antibody, VDw217 antibody, interleukin 17 receptor antibody

Format

0.5 mg IgG/ml in Tris saline (20mM Tris pH7.3, 150mM NaCl), 0.02% sodium azide, with 0.5% bovine serum albumin

Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

Goat Anti-IL17ra (mouse) Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

Goat Anti-IL17ra (mouse) Antibody - Protein Information**Goat Anti-IL17ra (mouse) Antibody - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)

- [Flow Cytometry](#)
- [Cell Culture](#)

Goat Anti-IL17ra (mouse) Antibody - Images



EB7882 (1 µg/ml) staining of Rat Kidney lysate (35 µg protein in RIPA buffer). Primary incubation was 1 hour. Detected by chemiluminescence.

Goat Anti-IL17ra (mouse) Antibody - References

Proinflammatory T helper type 17 cells are effective B-cell helpers. Mitsdoerffer M, et al. Proc Natl Acad Sci U S A, 2010 Aug 10. PMID 20660725.

Gut-residing segmented filamentous bacteria drive autoimmune arthritis via T helper 17 cells. Wu HJ, et al. Immunity, 2010 Jun 25. PMID 20620945.

Interleukin 17 receptor signaling is deleterious during Toxoplasma gondii infection in susceptible BL6 mice. Guiton R, et al. J Infect Dis, 2010 Aug 15. PMID 20575661.

Stem cell mobilization with G-CSF induces type 17 differentiation and promotes scleroderma. Hill GR, et al. Blood, 2010 Aug 5. PMID 20435882.

IL-17A-producing gammadelta T and Th17 lymphocytes mediate lung inflammation but not fibrosis in experimental silicosis. Lo Re S, et al. J Immunol, 2010 Jun 1. PMID 20421647.