

NOD1 / CARD4 Antibody (C-Term)
Peptide-affinity purified goat antibody
Catalog # AF2259a**Specification**

NOD1 / CARD4 Antibody (C-Term) - Product Information

Application	WB, E
Primary Accession	O9Y239
Other Accession	NP_006083.1 , 10392
Reactivity	Human
Predicted	Mouse, Rat, Pig, Dog
Host	Goat
Clonality	Polyclonal
Concentration	0.5 mg/ml
Isotype	IgG
Calculated MW	107691

NOD1 / CARD4 Antibody (C-Term) - Additional Information**Gene ID** 10392**Other Names**

Nucleotide-binding oligomerization domain-containing protein 1, Caspase recruitment domain-containing protein 4, NOD1, CARD4

Dilution

WB~~1:1000

E~~N/A

Format

0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin

Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

NOD1 / CARD4 Antibody (C-Term) is for research use only and not for use in diagnostic or therapeutic procedures.

NOD1 / CARD4 Antibody (C-Term) - Protein Information**Name** NOD1 {ECO:0000303|PubMed:10329646, ECO:0000312|HGNC:HGNC:16390}**Function**

Pattern recognition receptor (PRR) that detects bacterial peptidoglycan fragments and other danger signals and thus participates in both innate and adaptive immune responses (PubMed:<a

[11058605](http://www.uniprot.org/citations/11058605), PubMed: [12791997](http://www.uniprot.org/citations/12791997), PubMed: [12796777](http://www.uniprot.org/citations/12796777), PubMed: [15044951](http://www.uniprot.org/citations/15044951), PubMed: [16172124](http://www.uniprot.org/citations/16172124), PubMed: [19043560](http://www.uniprot.org/citations/19043560), PubMed: [22672233](http://www.uniprot.org/citations/22672233), PubMed: [27099311](http://www.uniprot.org/citations/27099311)). Specifically recognizes and binds gamma-D-glutamyl-meso-diaminopimelic acid (iE- DAP), a dipeptide present in peptidoglycan of Gram-negative bacteria (PubMed: [12791997](http://www.uniprot.org/citations/12791997), PubMed: [12796777](http://www.uniprot.org/citations/12796777), PubMed: [12871942](http://www.uniprot.org/citations/12871942), PubMed: [16172124](http://www.uniprot.org/citations/16172124), PubMed: [16211083](http://www.uniprot.org/citations/16211083)). Preferentially binds iE-DAP in tripeptide-containing muropetides (MurNAc-TriDAP or TriDAP) (PubMed: [16211083](http://www.uniprot.org/citations/16211083)). Ligand binding triggers oligomerization that facilitates the binding and subsequent activation of the proximal adapter receptor-interacting RIPK2 (PubMed: [12791997](http://www.uniprot.org/citations/12791997), PubMed: [12796777](http://www.uniprot.org/citations/12796777), PubMed: [17054981](http://www.uniprot.org/citations/17054981)). Following recruitment, RIPK2 undergoes 'Met-1'- (linear) and 'Lys-63'-linked polyubiquitination by E3 ubiquitin-protein ligases XIAP, BIRC2, BIRC3 and the LUBAC complex, becoming a scaffolding protein for downstream effectors, triggering activation of the NF-kappa-B and MAP kinases signaling (PubMed: [10880512](http://www.uniprot.org/citations/10880512), PubMed: [12791997](http://www.uniprot.org/citations/12791997), PubMed: [19043560](http://www.uniprot.org/citations/19043560)). This in turn leads to the transcriptional activation of hundreds of genes involved in immune response (PubMed: [10880512](http://www.uniprot.org/citations/10880512), PubMed: [19043560](http://www.uniprot.org/citations/19043560)). Also acts as a regulator of antiviral response elicited by dsRNA and the expression of RLR pathway members by targeting IFIH1 and TRAF3 to modulate the formation of IFIH1-MAVS and TRAF3-MAVS complexes leading to increased transcription of type I IFNs (PubMed: [32169843](http://www.uniprot.org/citations/32169843)). Also acts as a regulator of autophagy via its interaction with ATG16L1, possibly by recruiting ATG16L1 at the site of bacterial entry (By similarity). Besides recognizing pathogens, also involved in the endoplasmic reticulum stress response: acts by sensing and binding to the cytosolic metabolite sphingosine-1-phosphate generated in response to endoplasmic reticulum stress, initiating an inflammation process that leads to activation of the NF-kappa-B and MAP kinases signaling (PubMed: [27007849](http://www.uniprot.org/citations/27007849), PubMed: [33942347](http://www.uniprot.org/citations/33942347)). In addition, plays a role in insulin trafficking in beta cells in a cell-autonomous manner (By similarity). Mechanistically, upon recognizing cognate ligands, NOD1 and RIPK2 localize to insulin vesicles where they recruit RAB1A to direct insulin trafficking through the cytoplasm (By similarity).

Cellular Location

Cell membrane; Lipid-anchor. Apical cell membrane. Basolateral cell membrane. Cytoplasm. Note=Detected in the cytoplasm and at the cell membrane (PubMed:31649195). Following bacterial infection, localizes to bacterial entry sites in the cell membrane (PubMed:31649195). Recruited to the basolateral and apical membranes in polarized epithelial cells (PubMed:19043560)

Tissue Location

Highly expressed in adult heart, skeletal muscle, pancreas, spleen and ovary (PubMed:10224040).

Also detected in placenta, lung, liver, kidney, thymus, testis, small intestine and colon (PubMed:10224040).

NOD1 / CARD4 Antibody (C-Term) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

NOD1 / CARD4 Antibody (C-Term) - Images



AF2259a (0.5 µg/ml) staining of Human Duodenum lysate (35 µg protein in RIPA buffer). Primary incubation was 1 hour. Detected by chemiluminescence.

NOD1 / CARD4 Antibody (C-Term) - References

Association of NOD1 (CARD4) insertion/deletion polymorphism with susceptibility to IBD: a meta-analysis. Lu WG, Zou YF, Feng XL, Yuan FL, Gu YL, Li X, Li CW, Jin C, Li JP. World J Gastroenterol. 2010 Sep 14;16(34):4348-56. PMID: 20818820