

HDAC6 Antibody (C-Term)
Peptide-affinity purified goat antibody
Catalog # AF2262a**Specification**

HDAC6 Antibody (C-Term) - Product Information

Application	WB, E
Primary Accession	O9UBN7
Other Accession	NP_006035.2 , 10013
Reactivity	Human, Mouse
Predicted	Rat, Dog
Host	Goat
Clonality	Polyclonal
Concentration	0.5 mg/ml
Isotype	IgG
Calculated MW	131419

HDAC6 Antibody (C-Term) - Additional Information**Gene ID** 10013**Other Names**

Histone deacetylase 6, HD6, 3.5.1.98, HDAC6, KIAA0901

Dilution

WB~~1:1000

E~~N/A

Format

0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin

Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

HDAC6 Antibody (C-Term) is for research use only and not for use in diagnostic or therapeutic procedures.

HDAC6 Antibody (C-Term) - Protein Information**Name** HDAC6 {ECO:0000303|PubMed:10220385, ECO:0000312|HGNC:HGNC:14064}**Function**

Deacetylates a wide range of non-histone substrates (PubMed:12024216, PubMed:18606987, PubMed:18606987, PubMed:18606987)

[20308065](http://www.uniprot.org/citations/20308065), PubMed: [24882211](http://www.uniprot.org/citations/24882211), PubMed: [26246421](http://www.uniprot.org/citations/26246421), PubMed: [30538141](http://www.uniprot.org/citations/30538141), PubMed: [31857589](http://www.uniprot.org/citations/31857589), PubMed: [30770470](http://www.uniprot.org/citations/30770470), PubMed: [38534334](http://www.uniprot.org/citations/38534334), PubMed: [39567688](http://www.uniprot.org/citations/39567688)). Plays a central role in microtubule- dependent cell motility by mediating deacetylation of tubulin (PubMed: [12024216](http://www.uniprot.org/citations/12024216), PubMed: [20308065](http://www.uniprot.org/citations/20308065), PubMed: [26246421](http://www.uniprot.org/citations/26246421)). Required for cilia disassembly via deacetylation of alpha-tubulin (PubMed: [17604723](http://www.uniprot.org/citations/17604723), PubMed: [26246421](http://www.uniprot.org/citations/26246421)). Alpha-tubulin deacetylation results in destabilization of dynamic microtubules (By similarity). Promotes deacetylation of CTTN, leading to actin polymerization, promotion of autophagosome-lysosome fusion and completion of autophagy (PubMed: [30538141](http://www.uniprot.org/citations/30538141)). Deacetylates SQSTM1 (PubMed: [31857589](http://www.uniprot.org/citations/31857589) target="_blank">31857589). Deacetylates peroxiredoxins PRDX1 and PRDX2, decreasing their reducing activity (PubMed: [18606987](http://www.uniprot.org/citations/18606987) target="_blank">18606987). Deacetylates antiviral protein RIGI in the presence of viral mRNAs which is required for viral RNA detection by RIGI (By similarity). Sequentially deacetylates and polyubiquitinates DNA mismatch repair protein MSH2 which leads to MSH2 degradation, reducing cellular sensitivity to DNA-damaging agents and decreasing cellular DNA mismatch repair activities (PubMed: [24882211](http://www.uniprot.org/citations/24882211) target="_blank">24882211). Deacetylates DNA mismatch repair protein MLH1 which prevents recruitment of the MutL alpha complex (formed by the MLH1-PMS2 heterodimer) to the MutS alpha complex (formed by the MSH2-MSH6 heterodimer), leading to tolerance of DNA damage (PubMed: [30770470](http://www.uniprot.org/citations/30770470) target="_blank">30770470). Deacetylates RHOT1/MIRO1 which blocks mitochondrial transport and mediates axon growth inhibition (By similarity). Deacetylates transcription factor SP1 which leads to increased expression of ENG, positively regulating angiogenesis (PubMed: [38534334](http://www.uniprot.org/citations/38534334) target="_blank">38534334). Deacetylates KHDRBS1/SAM68 which regulates alternative splicing by inhibiting the inclusion of CD44 alternate exons (PubMed: [26080397](http://www.uniprot.org/citations/26080397) target="_blank">26080397). Acts as a valine sensor by binding to valine through the primate-specific SE14 repeat region (PubMed: [39567688](http://www.uniprot.org/citations/39567688) target="_blank">39567688). In valine deprivation conditions, translocates from the cytoplasm to the nucleus where it deacetylates TET2 which promotes TET2-dependent DNA demethylation, leading to DNA damage (PubMed: [39567688](http://www.uniprot.org/citations/39567688) target="_blank">39567688). Promotes odontoblast differentiation following IPO7-mediated nuclear import and subsequent repression of RUNX2 expression (By similarity). In addition to its protein deacetylase activity, plays a key role in the degradation of misfolded proteins: when misfolded proteins are too abundant to be degraded by the chaperone refolding system and the ubiquitin-proteasome, mediates the transport of misfolded proteins to a cytoplasmic juxtanuclear structure called aggresome (PubMed: [17846173](http://www.uniprot.org/citations/17846173) target="_blank">17846173). Probably acts as an adapter that recognizes polyubiquitinated misfolded proteins and targets them to the aggresome, facilitating their clearance by autophagy (PubMed: [17846173](http://www.uniprot.org/citations/17846173) target="_blank">17846173). Involved in the MTA1-mediated epigenetic regulation of ESR1 expression in breast cancer (PubMed: [24413532](http://www.uniprot.org/citations/24413532) target="_blank">24413532).

Cellular Location

Cytoplasm. Cytoplasm, cytoskeleton. Nucleus. Perikaryon {ECO:0000250|UniProtKB:Q9Z2V5}. Cell

projection, dendrite {ECO:0000250|UniProtKB:Q9Z2V5}. Cell projection, axon {ECO:0000250|UniProtKB:Q9Z2V5}. Cell projection, cilium. Cytoplasm, cytoskeleton, microtubule organizing center, centrosome. Cytoplasm, cytoskeleton, cilium basal body Note=Mainly cytoplasmic where it is associated with microtubules (PubMed:12024216). Can shuttle between the cytoplasm and the nucleus (PubMed:39567688). Retained in the cytoplasm by binding to valine via the primate-specific SE14 repeat region while valine deprivation induces nuclear localization (PubMed:39567688). Found exclusively in the cytoplasm in proliferative cells with a fraction found in the nucleus during differentiation (By similarity). May translocate to the nucleus following DNA damage (PubMed:30770470) {ECO:0000250|UniProtKB:Q9Z2V5, ECO:0000269|PubMed:12024216, ECO:0000269|PubMed:30770470, ECO:0000269|PubMed:39567688}

HDAC6 Antibody (C-Term) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

HDAC6 Antibody (C-Term) - Images



AF2262a (1 µg/ml) staining of HeLa nuclear lysate (35 µg protein in RIPA buffer). Primary incubation was 1 hour. Detected by chemiluminescence.

HDAC6 Antibody (C-Term) - References

HDAC6 modulates cell motility by altering the acetylation level of cortactin. Zhang X, Yuan Z, Zhang Y, Yong S, Salas-Burgos A, Koomen J, Olashaw N, Parsons JT, Yang XJ, Dent SR, Yao TP, Lane WS, Seto E. Mol Cell. 2007 Jul 20;27(2):197-213 PMID: 17643370