

**RANBP8 / IPO8 Antibody (C-Term)**  
**Peptide-affinity purified goat antibody**  
**Catalog # AF2272a****Specification**

---

**RANBP8 / IPO8 Antibody (C-Term) - Product Information**

Application	IHC, E
Primary Accession	<a href="#">O15397</a>
Other Accession	<a href="#">NP_006381.2</a> , <a href="#">NP_001177924.1</a> , <a href="#">10526</a>
Reactivity	Human
Predicted	Pig, Dog
Host	Goat
Clonality	Polyclonal
Concentration	0.5 mg/ml
Isotype	IgG
Calculated MW	119938

**RANBP8 / IPO8 Antibody (C-Term) - Additional Information****Gene ID** 10526**Other Names**

Importin-8, Imp8, Ran-binding protein 8, RanBP8, IPO8, RANBP8

**Dilution**

IHC~~1:100~500

E~~N/A

**Format**

0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin

**Storage**

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

**Precautions**

RANBP8 / IPO8 Antibody (C-Term) is for research use only and not for use in diagnostic or therapeutic procedures.

**RANBP8 / IPO8 Antibody (C-Term) - Protein Information****Name** IPO8**Synonyms** RANBP8**Function**

Involved in nuclear protein import, either by acting as autonomous nuclear transport receptor or

as an adapter-like protein in association with the importin-beta subunit KPNB1. Acting autonomously, may serve as receptor for nuclear localization signals (NLS) and promote translocation of import substrates through the nuclear pore complex (NPC) by an energy requiring, Ran-dependent mechanism. At the nucleoplasmic side of the NPC, Ran binds to importin, the importin/substrate complex dissociates and importin is re-exported from the nucleus to the cytoplasm where GTP hydrolysis releases Ran. The directionality of nuclear import is thought to be conferred by an asymmetric distribution of the GTP- and GDP-bound forms of Ran between the cytoplasm and nucleus (PubMed:<a href="http://www.uniprot.org/citations/9214382" target="\_blank">9214382</a>). In vitro mediates the nuclear import of the signal recognition particle protein SRP19 (PubMed:<a href="http://www.uniprot.org/citations/11682607" target="\_blank">11682607</a>). May also be involved in cytoplasm-to-nucleus shuttling of a broad spectrum of other cargos, including Argonaute- microRNAs complexes, the JUN protein, RELA/NF-kappa-B p65 subunit, the translation initiation factor EIF4E and a set of receptor-activated mothers against decapentaplegic homolog (SMAD) transcription factors that play a critical role downstream of the large family of transforming growth factor beta and bone morphogenetic protein (BMP) cytokines (Probable).

#### **Cellular Location**


Cytoplasm. Nucleus.

#### **RANBP8 / IPO8 Antibody (C-Term) - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

#### **RANBP8 / IPO8 Antibody (C-Term) - Images**



AF2272a (4 µg/ml) staining of paraffin embedded Human Tonsil. Steamed antigen retrieval with Tris/EDTA buffer pH 9, HRP-staining. Similar results were obtained after antigen retrieval at pH9.

#### **RANBP8 / IPO8 Antibody (C-Term) - Background**

This antibody is expected to recognise isoform 1 (NP\_006381.2) and isoform 2 (NP\_001177924.1).

**RANBP8 / IPO8 Antibody (C-Term) - References**

A novel class of RanGTP binding proteins. Gorlich D, Dabrowski M, Bischoff FR, Kutay U, Bork P, Hartmann E, Prehn S, Izaurralde E. J Cell Biol. 1997 Jul 14;138(1):65-80. PMID: 9214382