

### CBX6 and CBX8 Antibody (N-Term)

Peptide-affinity purified goat antibody Catalog # AF2306a

## **Specification**

## CBX6 and CBX8 Antibody (N-Term) - Product Information

Application

Primary Accession <u>095503</u>

Other Accession <u>NP\_055107.3</u>, <u>NP\_065700.1</u>, <u>23466</u>, <u>57332</u>

Predicted Human, Mouse, Rat, Pig, Dog

Host Goat
Clonality Polyclonal
Concentration 0.5 mg/ml
Isotype IgG

Isotype IgG
Calculated MW 43898

# CBX6 and CBX8 Antibody (N-Term) - Additional Information

## **Gene ID 23466**

### **Other Names**

Chromobox protein homolog 6, CBX6

## **Dilution**

E~~N/A

#### **Format**

0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin

## **Storage**

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

#### **Precautions**

CBX6 and CBX8 Antibody (N-Term) is for research use only and not for use in diagnostic or therapeutic procedures.

# CBX6 and CBX8 Antibody (N-Term) - Protein Information

#### Name CBX6

#### **Function**

Component of a Polycomb group (PcG) multiprotein PRC1-like complex, a complex class required to maintain the transcriptionally repressive state of many genes, including Hox genes, throughout development (PubMed:<a href="http://www.uniprot.org/citations/21282530" target="\_blank">21282530</a>). PcG PRC1 complex acts via chromatin remodeling and modification of histones; it mediates monoubiquitination of histone H2A 'Lys-119', rendering



Tel: 858.875.1900 Fax: 858.875.1999

chromatin heritably changed in its expressibility. Possibly contributes to the target selectivity of the PRC1 complex by binding specific regions of chromatin (PubMed: <a href="http://www.uniprot.org/citations/18927235" target=" blank">18927235</a>). Recruitment to chromatin might occur in an H3K27me3- independent fashion (By similarity). May have a PRC1-independent function in embryonic stem cells (By similarity).

#### **Cellular Location**

Nucleus. Chromosome. Note=Uniformely distributed in the nucleoplasm (PubMed:18927235). Localizes to the inactivated X chromosome in females (By similarity). {ECO:0000250|UniProtKB:Q9DBY5, ECO:0000269|PubMed:18927235}

## CBX6 and CBX8 Antibody (N-Term) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

### CBX6 and CBX8 Antibody (N-Term) - Images

# CBX6 and CBX8 Antibody (N-Term) - Background

This antibody is expected to recognise both the human proteins CBX6 and CBX8, which are almost identical.

# CBX6 and CBX8 Antibody (N-Term) - References

HPC3 is a new human polycomb orthologue that interacts and associates with RING1 and Bmi1 and has transcriptional repression properties. Bardos JI, Saurin AJ, Tissot C, Duprez E, Freemont PS. | Biol Chem. 2000 Sep 15;275(37):28785-92. PMID: 10825164