

SSA1 / Ro52 Antibody (C-Term)

Peptide-affinity purified goat antibody Catalog # AF2325a

Specification

SSA1 / Ro52 Antibody (C-Term) - Product Information

Application WB, E
Primary Accession P19474

Other Accession NP 003132.2, 6737

Reactivity
Host
Clonality
Concentration

Human
Goat
Polyclonal
O.5 mg/ml

Isotype IgG
Calculated MW 54170

SSA1 / Ro52 Antibody (C-Term) - Additional Information

Gene ID 6737

Other Names

E3 ubiquitin-protein ligase TRIM21, 6.3.2.-, 52 kDa Ro protein, 52 kDa ribonucleoprotein autoantigen Ro/SS-A, RING finger protein 81, Ro(SS-A), Sjoegren syndrome type A antigen, SS-A, Tripartite motif-containing protein 21, TRIM21, RNF81, RO52, SSA1

Dilution

WB~~1:1000 E~~N/A

Format

0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin

Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

SSA1 / Ro52 Antibody (C-Term) is for research use only and not for use in diagnostic or therapeutic procedures.

SSA1 / Ro52 Antibody (C-Term) - Protein Information

Name TRIM21 (HGNC:11312)

Synonyms RNF81, RO52, SSA1

Function



and UBE2E2 (PubMed: 16297862, PubMed:16316627, PubMed:16472766, PubMed:16880511, PubMed:18022694, PubMed:18361920, PubMed:18641315, PubMed:18845142, PubMed:19675099, PubMed:26347139). Forms a ubiquitin ligase complex in cooperation with the E2 UBE2D2 that is used not only for the ubiquitination of USP4 and IKBKB but also for its self-ubiquitination (PubMed: 16880511, PubMed:19675099). Component of cullin-RING-based SCF (SKP1-CUL1-F-box protein) E3 ubiquitin-protein ligase complexes such as SCF(SKP2)-like complexes (PubMed: 16880511). A TRIM21-containing SCF(SKP2)-like complex is shown to mediate ubiquitination of CDKN1B ('Thr-187' phosphorylated- form), thereby promoting its degradation by the proteasome (PubMed:16880511). Monoubiquitinates IKBKB that will negatively regulates Tax-induced NF-kappa-B signaling (PubMed:19675099). Negatively regulates IFN-beta production post-pathogen recognition by catalyzing polyubiquitin-mediated degradation of IRF3 (PubMed:18641315). Mediates the ubiquitin-mediated proteasomal degradation of IgG1 heavy chain, which is linked to the VCP-mediated ER-associated degradation (ERAD) pathway (PubMed: 18022694). Promotes IRF8 ubiquitination, which enhanced the ability of IRF8 to stimulate cytokine genes transcription in macrophages (By similarity). Plays a role in the regulation of the cell cycle progression (PubMed:16880511). Enhances the decapping activity of DCP2 (PubMed:18361920). Exists as a ribonucleoprotein particle present in all mammalian cells studied and composed of a single polypeptide and one of four small RNA molecules (PubMed:1985094, PubMed:8666824). At least two isoforms are present in nucleated and red blood cells, and tissue specific differences in RO/SSA proteins have been identified (PubMed:8666824). The common feature of these proteins is their ability to bind HY RNAs.2 (PubMed:8666824). Involved in the regulation of innate immunity and the inflammatory response in response to IFNG/IFN-gamma (PubMed: 26347139). Organizes autophagic machinery by serving as a platform for the assembly of ULK1, Beclin 1/BECN1 and ATG8 family members and recognizes specific autophagy targets, thus coordinating target recognition with assembly of the autophagic apparatus and initiation of autophagy (PubMed: 26347139). Also regulates autophagy through FIP200/RB1CC1 ubiquitination and subsequent decreased protein stability (PubMed:36359729). Represses the innate antiviral response by facilitating the formation of the NMI-IFI35 complex through 'Lys-63'- linked ubiquitination of NMI (PubMed: 26342464). During viral infection, promotes cell pyroptosis by mediating 'Lys-6'-linked ubiquitination of ISG12a/IFI27, facilitating its translocation into the mitochondria and subsequent CASP3 activation (PubMed: <a $href="http://www.uniprot.org/citations/36426955"\ target="_blank">36426955).\ When$

E3 ubiquitin-protein ligase whose activity is dependent on E2 enzymes, UBE2D1, UBE2D2, UBE2E1



up-regulated through the IFN/JAK/STAT signaling pathway, promotes 'Lys-27'-linked ubiquitination of MAVS, leading to the recruitment of TBK1 and up- regulation of innate immunity (PubMed:29743353). Mediates 'Lys-63'- linked polyubiquitination of G3BP1 in response to heat shock, leading to stress granule disassembly (PubMed:36692217).

Cellular Location

Cytoplasm. Cytoplasmic vesicle, autophagosome. Nucleus. Cytoplasm, P-body. Cytoplasm, Stress granule. Note=Enters the nucleus upon exposure to nitric oxide (PubMed:18361920). Localizes to small dot- or rod-like structures in the cytoplasm, called processing bodies (P-bodies) that are located underneath the plasma membrane and also diffusely in the cytoplasm (PubMed:18361920). They are located along the microtubules and are highly motile in cells (PubMed:18361920). Colocalizes with DCP2 in P-bodies (PubMed:18361920). Localizes to stress granules in response to oxidative stress (PubMed:36692217).

Tissue Location


Isoform 1 and isoform 2 are expressed in fetal and adult heart and fetal lung

SSA1 / Ro52 Antibody (C-Term) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

SSA1 / Ro52 Antibody (C-Term) - Images



HEK293 lysate (10ug protein in RIPA buffer) overexpressing Human TRIM21 with C-terminal MYC tag probed with AF2325a (1ug/ml) in Lane A and probed with anti-MYC Tag (1/1000) in lane C. Mock-transfected HEK293 probed with AF2325a (1mg/ml) in Lane B. Primary incubations were for



Tel: 858.875.1900 Fax: 858.875.1999



1 hour. Detected by chemiluminescence.

SSA1 / Ro52 Antibody (C-Term) - References

52-kD SS-A/Ro: genomic structure and identification of an alternatively spliced transcript encoding a novel leucine zipper-minus autoantigen expressed in fetal and adult heart. Chan EK, Di Donato F, Hamel JC, Tseng CE, Buyon JP. J Exp Med. 1995 Oct 1;182(4):983-92. PMID: 7561701