

ECSM2 Antibody (C-Term)
Peptide-affinity purified goat antibody
Catalog # AF2410a**Specification**

ECSM2 Antibody (C-Term) - Product Information

Application	WB
Primary Accession	Q19T08.1
Other Accession	NP_001071161.1
Reactivity	Human
Host	Goat
Clonality	Polyclonal
Concentration	0.5 mg/ml
Isotype	IgG

ECSM2 Antibody (C-Term) - Additional Information**Other Names**

endothelial cell-specific marker 2, endothelial cell-specific molecule 2; hypothetical protein
LOC641700

Format

0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin

Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.


Precautions

ECSM2 Antibody (C-Term) is for research use only and not for use in diagnostic or therapeutic procedures.

ECSM2 Antibody (C-Term) - Protein Information**ECSM2 Antibody (C-Term) - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

ECSM2 Antibody (C-Term) - Images

AF2410a (1 µg/ml) staining of Human lung (A) and Human Umbical cord (B) lysates (35 µg protein in RIPA buffer). Primary incubation was 1 hour.

ECSM2 Antibody (C-Term) - References

PMID: 14970008