

RIF1 (aa 2076-2089) Antibody (internal region)
Peptide-affinity purified goat antibody
Catalog # AF2421a**Specification**

RIF1 (aa 2076-2089) Antibody (internal region) - Product Information

Application	E
Primary Accession	Q5UIPO
Other Accession	NP_060621.3 , 55183
Predicted	Human
Host	Goat
Clonality	Polyclonal
Concentration	0.5 mg/ml
Isotype	IgG
Calculated MW	274466

RIF1 (aa 2076-2089) Antibody (internal region) - Additional Information**Gene ID** 55183**Other Names**

Telomere-associated protein RIF1, Rap1-interacting factor 1 homolog, RIF1

Dilution

E~~N/A

Format

0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin

Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

RIF1 (aa 2076-2089) Antibody (internal region) is for research use only and not for use in diagnostic or therapeutic procedures.

RIF1 (aa 2076-2089) Antibody (internal region) - Protein Information**Name** RIF1 {ECO:0000303|PubMed:15342490, ECO:0000312|HGNC:HGNC:23207}**Function**

Key regulator of TP53BP1 that plays a key role in the repair of double-strand DNA breaks (DSBs) in response to DNA damage: acts by promoting non-homologous end joining (NHEJ)-mediated repair of DSBs (PubMed: <http://www.uniprot.org/citations/15342490> target="_blank">15342490, PubMed: <http://www.uniprot.org/citations/28241136> target="_blank">28241136). In response to DNA damage, interacts with ATM-phosphorylated

TP53BP1 (PubMed:23333306, PubMed:28241136). Interaction with TP53BP1 leads to dissociate the interaction between NUDT16L1/TIRR and TP53BP1, thereby unmasking the tandem Tudor-like domain of TP53BP1 and allowing recruitment to DNA DSBs (PubMed:28241136). Once recruited to DSBs, RIF1 and TP53BP1 act by promoting NHEJ-mediated repair of DSBs (PubMed:23333306). In the same time, RIF1 and TP53BP1 specifically counteract the function of BRCA1 by blocking DSBs resection via homologous recombination (HR) during G1 phase (PubMed:23333306). Also required for immunoglobulin class-switch recombination (CSR) during antibody genesis, a process that involves the generation of DNA DSBs (By similarity). Promotes NHEJ of dysfunctional telomeres (By similarity).

Cellular Location

Nucleus. Chromosome {ECO:0000250|UniProtKB:Q6PR54}. Chromosome, telomere. Cytoplasm, cytoskeleton, spindle. Note=Following interaction with TP53BP1, recruited to sites of DNA damage, such as DSBs (By similarity). Exhibits ATM- and TP53BP1-dependent localization to uncapped or aberrant telomeres and to DNA double strand breaks (DSBs) (PubMed:15342490). Does not associate with normal telomere structures (PubMed:15342490, PubMed:15583028). Localizes to microtubules of the midzone of the mitotic spindle during anaphase, and to condensed chromosomes in telophase (PubMed:15583028) {ECO:0000250|UniProtKB:Q6PR54, ECO:0000269|PubMed:15342490, ECO:0000269|PubMed:15583028}

Tissue Location

Highly expressed in testis.

RIF1 (aa 2076-2089) Antibody (internal region) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

RIF1 (aa 2076-2089) Antibody (internal region) - Images

RIF1 (aa 2076-2089) Antibody (internal region) - Background

This antibody is expected to recognise an epitope corresponding to aa 2076-2089 of human RIF1 protein.

RIF1 (aa 2076-2089) Antibody (internal region) - References

Human Rif1, ortholog of a yeast telomeric protein, is regulated by ATM and 53BP1 and functions in the S-phase checkpoint. Silverman J, Takai H, Buonomo SB, Eisenhaber F, de Lange T. Genes Dev. 2004 Sep 1;18(17):2108-19. PMID: 15342490