

NEUROD2 Antibody (C-Term)
Peptide-affinity purified goat antibody
Catalog # AF2492a**Specification**

NEUROD2 Antibody (C-Term) - Product Information

Application	E
Primary Accession	Q15784
Other Accession	NP_006151.3 , 4761
Predicted	Human, Mouse, Rat, Dog, Cow
Host	Goat
Clonality	Polyclonal
Concentration	0.5 mg/ml
Isotype	IgG
Calculated MW	41361

NEUROD2 Antibody (C-Term) - Additional Information**Gene ID** 4761**Other Names**

Neurogenic differentiation factor 2, NeuroD2, Class A basic helix-loop-helix protein 1, bHLHa1, NeuroD-related factor, NDRF, NEUROD2, BHLHA1, NDRF

Format

0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin

Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

NEUROD2 Antibody (C-Term) is for research use only and not for use in diagnostic or therapeutic procedures.

NEUROD2 Antibody (C-Term) - Protein Information**Name** NEUROD2**Synonyms** BHLHA1, NDRF**Function**

Transcriptional regulator implicated in neuronal determination. Mediates calcium-dependent transcription activation by binding to E box-containing promoter. Critical factor essential for the repression of the genetic program for neuronal differentiation; prevents the formation of synaptic vesicle clustering at active zone to the presynaptic membrane in postmitotic neurons. Induces transcription of ZEB1, which in turn represses neuronal differentiation by down- regulating REST

expression. Plays a role in the establishment and maturation of thalamocortical connections; involved in the segregation of thalamic afferents into distinct barrel domains within layer VI of the somatosensory cortex. Involved in the development of the cerebellar and hippocampal granular neurons, neurons in the basolateral nucleus of amygdala and the hypothalamic-pituitary axis. Associates with chromatin to the DPYSL3 E box-containing promoter (By similarity).

Cellular Location

Nucleus {ECO:0000255|PROSITE-ProRule:PRU00981}.

NEUROD2 Antibody (C-Term) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

NEUROD2 Antibody (C-Term) - Images**NEUROD2 Antibody (C-Term) - References**

The dosage of the neuroD2 transcription factor regulates amygdala development and emotional learning. Lin CH, Hansen S, Wang Z, Storm DR, Tapscott SJ, Olson JM. Proc Natl Acad Sci U S A. 2005 Oct 11;102(41):14877-82. Epub 2005 Oct 3. PMID: 16203979