

STELLAR Antibody (internal region)
Peptide-affinity purified goat antibody
Catalog # AF2502a**Specification**

STELLAR Antibody (internal region) - Product Information

Application	Pep-ELISA
Primary Accession	O6W0C5.1
Other Accession	NP_954980.1 , 400206
Predicted	Human
Host	Goat
Clonality	Polyclonal
Concentration	0.5 mg/ml
Isotype	IgG

STELLAR Antibody (internal region) - Additional Information**Other Names**

STELLAR; germ and embryonic stem cell enriched protein STELLA ; developmental pluripotency associated 3 pseudogene, stella-like; STELLA;

Format

0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin

Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

STELLAR Antibody (internal region) is for research use only and not for use in diagnostic or therapeutic procedures.

STELLAR Antibody (internal region) - Protein Information**STELLAR Antibody (internal region) - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

STELLAR Antibody (internal region) - Images**STELLAR Antibody (internal region) - References**

Human embryonic stem cell genes OCT4, NANOG, STELLAR, and GDF3 are expressed in both seminoma and breast carcinoma. Ezech UI, Turek PJ, Reijo RA, Clark AT. Cancer. 2005 Nov 15;104(10):2255-65. PMID: 16228988