

## PGRMC2 Antibody (C-Term)

Peptide-affinity purified goat antibody Catalog # AF2514a

## **Specification**

## PGRMC2 Antibody (C-Term) - Product Information

Application

Primary Accession <u>O15173</u>

Other Accession NP 006311.2, 10424

Predicted Human, Mouse, Rat, Pig, Dog, Cow

Host Goat
Clonality Polyclonal
Concentration 0.5 mg/ml

Isotype IgG
Calculated MW 23818

# PGRMC2 Antibody (C-Term) - Additional Information

#### **Gene ID 10424**

## **Other Names**

Membrane-associated progesterone receptor component 2, Progesterone membrane-binding protein, Steroid receptor protein DG6, PGRMC2, DG6, PMBP

#### **Format**

0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin

## Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

### **Precautions**

PGRMC2 Antibody (C-Term) is for research use only and not for use in diagnostic or therapeutic procedures.

# PGRMC2 Antibody (C-Term) - Protein Information

## Name PGRMC2 (HGNC:16089)

#### **Function**

Required for the maintenance of uterine histoarchitecture and normal female reproductive lifespan (By similarity). May serve as a universal non-classical progesterone receptor in the uterus (Probable). Intracellular heme chaperone required for delivery of labile, or signaling heme, to the nucleus (By similarity). Plays a role in adipocyte function and systemic glucose homeostasis (PubMed:<a href="http://www.uniprot.org/citations/28111073" target="\_blank">28111073</a>). In brown fat, which has a high demand for heme, delivery of labile heme in the nucleus regulates the activity of heme-responsive transcriptional repressors such as NR1D1 and BACH1 (By



Tel: 858.875.1900 Fax: 858.875.1999

similarity).

### **Cellular Location**

Membrane; Single- pass membrane protein. Nucleus envelope. Endoplasmic reticulum. Secreted

### **Tissue Location**

Expressed by endometrial glands and stroma (at protein level) (PubMed:23793472). Detected in urine (at protein level) (PubMed:37453717).

## **PGRMC2 Antibody (C-Term) - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- <u>Immunohistochemistry</u>
- Immunofluorescence
- <u>Immunoprecipitation</u>
- Flow Cytomety
- Cell Culture

# PGRMC2 Antibody (C-Term) - Images

## PGRMC2 Antibody (C-Term) - References

Placental site trophoblastic tumor: human placental lactogen and pregnancy-associated major basic protein as immunohistologic markers. Rhoton-Vlasak A, Wagner JM, Rutgers JL, Baergen RN, Young RH, Roche PC, Plummer TB, Gleich GJ. Hum Pathol. 1998 Mar;29(3):280-8. PMID: 9496832