

Timp2 (zebrafish) Antibody (internal region)
Peptide-affinity purified goat antibody
Catalog # AF2609a**Specification**

Timp2 (zebrafish) Antibody (internal region) - Product Information

Application	E
Other Accession	NP_878294.1
Predicted	Zebrafish
Host	Goat
Clonality	Polyclonal
Concentration	0.5 mg/ml
Isotype	IgG

Timp2 (zebrafish) Antibody (internal region) - Additional Information**Other Names**

timp2; tissue inhibitor of metalloproteinase 2 ; MGC136726; etID32613.12; fa96h09; wu:fa96h09; zgc:136726

Format

0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin

Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

Timp2 (zebrafish) Antibody (internal region) is for research use only and not for use in diagnostic or therapeutic procedures.

Timp2 (zebrafish) Antibody (internal region) - Protein Information**Timp2 (zebrafish) Antibody (internal region) - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Timp2 (zebrafish) Antibody (internal region) - Images

Timp2 (zebrafish) Antibody (internal region) - Background

This antibody is not expected to cross-react to Timp2L.

Timp2 (zebrafish) Antibody (internal region) - References

The expression of tissue inhibitor of metalloproteinase 2 (TIMP-2) is required for normal development of zebrafish embryos. Zhang J, Bai S, Tanase C, Nagase H, Sarras MP Jr. Dev Genes Evol. 2003 Aug;213(8):382-9. Epub 2003 May 8. PMID: 12736828