

Cezanne2 Antibody (internal region)
Peptide-affinity purified goat antibody
Catalog # AF2796a**Specification**

Cezanne2 Antibody (internal region) - Product Information

Application	E
Primary Accession	Q8TE49
Other Accession	NP_570971.1 , 161725 , 170711 (mouse)
Predicted	Human, Mouse
Host	Goat
Clonality	Polyclonal
Concentration	0.5 mg/ml
Isotype	IgG
Calculated MW	100677

Cezanne2 Antibody (internal region) - Additional Information**Gene ID** 161725**Other Names**

OTU domain-containing protein 7A, 3.4.19.12, Zinc finger protein Cezanne 2, OTUD7A, C15orf16, CEZANNE2, OTUD7

Dilution

E~~N/A

Format

0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin

Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

Cezanne2 Antibody (internal region) is for research use only and not for use in diagnostic or therapeutic procedures.

Cezanne2 Antibody (internal region) - Protein Information**Name** OTUD7A**Synonyms** C15orf16, CEZANNE2, OTUD7**Function**

Deubiquitinase, which cleaves 'Lys-11'-linked polyubiquitin chains. Might be required for PA28-20S proteasome assembly (Probable).

Cellular Location

Cytoplasm. Nucleus.

Cezanne2 Antibody (internal region) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Cezanne2 Antibody (internal region) - Images**Cezanne2 Antibody (internal region) - References**

Differential gene expression of TRPM1, the potential cause of congenital stationary night blindness and coat spotting patterns (LP) in the Appaloosa horse (*Equus caballus*). Bellone RR, Brooks SA, Sandmeyer L, Murphy BA, Forsyth G, Archer S, Bailey E, Grahn B. Genetics. 2008 Aug;179(4):1861-70. Epub 2008 Jul 27. PMID: 18660533