

CLMP Antibody (C-Term)

Peptide-affinity purified goat antibody Catalog # AF2845a

Specification

CLMP Antibody (C-Term) - Product Information

Application Primary Accession Other Accession Reactivity Host Clonality Concentration Isotype Calculated MW WB, E <u>Q9H6B4</u> <u>NP_079045.1</u>, <u>79827</u> Human Goat Polyclonal 0.5 mg/ml lgG 41281

CLMP Antibody (C-Term) - Additional Information

Gene ID 79827

Other Names

CXADR-like membrane protein, Adipocyte adhesion molecule, Coxsackie- and adenovirus receptor-like membrane protein, CAR-like membrane protein, CLMP, ACAM, ASAM

Dilution WB~~1:1000 E~~N/A

Format 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin

Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

CLMP Antibody (C-Term) is for research use only and not for use in diagnostic or therapeutic procedures.

CLMP Antibody (C-Term) - Protein Information

Name CLMP

Synonyms ACAM, ASAM

Function

May be involved in the cell-cell adhesion. May play a role in adipocyte differentiation and



development of obesity. Is required for normal small intestine development.

Cellular Location

Cell junction, tight junction. Cell membrane; Single-pass type I membrane protein

Tissue Location

Predominantly expressed in epithelial cells within different tissues and in the white adipose tissue. Expressed at high levels in small intestine and placenta, at intermediate levels in the heart, skeletal muscle, colon, spleen, kidney and lung and at low levels in the liver and peripheral blood leukocytes. Highly abundant in the intestine during embryo and fetal development (at protein level)

CLMP Antibody (C-Term) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- <u>Western Blot</u>
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- <u>Cell Culture</u>

CLMP Antibody (C-Term) - Images



AF2845a staining (0.5 μ g/ml) of Human Placenta lysate (RIPA buffer, 35 μ g total protein per lane). Primary incubated for 1 hour. Detected by western blot using chemiluminescence.

CLMP Antibody (C-Term) - References

CLMP, a novel member of the CTX family and a new component of epithelial tight junctions. Raschperger E, Engstrom U, Pettersson RF, Fuxe J. J Biol Chem. 2004 Jan 2;279(1):796-804. PMID: 14573622