

ITM2B Antibody (internal region)
Peptide-affinity purified goat antibody
Catalog # AF2903a**Specification**

ITM2B Antibody (internal region) - Product Information

Application	WB, E
Primary Accession	Q9Y287
Other Accession	NP_068839.1 , 9445 , 16432 (mouse) , 290364 (rat)
Reactivity	Human
Predicted	Mouse, Rat
Host	Goat
Clonality	Polyclonal
Concentration	0.5 mg/ml
Isotype	IgG
Calculated MW	30338

ITM2B Antibody (internal region) - Additional Information**Gene ID** 9445**Other Names**

Integral membrane protein 2B, Immature BRI2, imBRI2, Protein E25B, Transmembrane protein BRI, Bri, BRI2, membrane form, Mature BRI2, mBRI2, BRI2 intracellular domain, BRI2 ICD, BRI2C, soluble form, Bri23 peptide, Bri2-23, ABri23, C-terminal peptide, P23 peptide, ITM2B, BRI, BRI2

Dilution

WB~~1:1000

E~~N/A

Format

0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin

Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

ITM2B Antibody (internal region) is for research use only and not for use in diagnostic or therapeutic procedures.

ITM2B Antibody (internal region) - Protein Information**Name** ITM2B**Synonyms** BRI, BRI2

Function

Plays a regulatory role in the processing of the amyloid-beta A4 precursor protein (APP) and acts as an inhibitor of the amyloid-beta peptide aggregation and fibrils deposition. Plays a role in the induction of neurite outgrowth. Functions as a protease inhibitor by blocking access of secretases to APP cleavage sites. Bri23 peptide prevents aggregation of APP amyloid-beta protein 42 into toxic oligomers.

Cellular Location

[Integral membrane protein 2B]: Golgi apparatus membrane; Single-pass type II membrane protein Note=Immature BRI2 (imBRI2) is cleaved by furin in the Golgi into mBRI2 and a Bri23 peptide. mBRI2 is transported to the plasma membrane and Bri23 peptide is secreted [Bri23 peptide]: Secreted. Note=Detected in the cerebral spinal fluid (CSF).

Tissue Location

Ubiquitous. Expressed in brain.

ITM2B Antibody (internal region) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

ITM2B Antibody (internal region) - Images

AF2903a (1 µg/ml) staining of Human Hippocampus lysate (35 µg protein in RIPA buffer). Primary incubation was 1 hour. Detected by chemiluminescence.

ITM2B Antibody (internal region) - References

Understanding the development of human bladder cancer by using a whole-organ genomic mapping strategy. Majewski T, Lee S, Jeong J, Yoon DS, Kram A, Kim MS, Tuziak T, Bondaruk J, Lee S, Park WS, Tang KS, Chung W, Shen L, Ahmed SS, Johnston DA, Grossman HB, Dinney CP, Zhou JH, Harris RA, Snyder C, Filipek S, Narod SA, Watson P, Lynch HT, Gazdar A, Bar-Eli M, Lab. Invest. 2008 Jul 88 (7): 694-721. PMID: 18458673