

SAFB2 Antibody (internal region)
Peptide-affinity purified goat antibody
Catalog # AF2937a**Specification**

SAFB2 Antibody (internal region) - Product Information

Application	E
Primary Accession	Q14151
Other Accession	NP_055464.1 , 9667
Predicted	Human
Host	Goat
Clonality	Polyclonal
Concentration	0.5 mg/ml
Isotype	IgG
Calculated MW	107473

SAFB2 Antibody (internal region) - Additional Information**Gene ID** 9667**Other Names**

Scaffold attachment factor B2, SAF-B2, SAFB2, KIAA0138

Format

0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin

Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

SAFB2 Antibody (internal region) is for research use only and not for use in diagnostic or therapeutic procedures.

SAFB2 Antibody (internal region) - Protein Information**Name** SAFB2**Synonyms** KIAA0138**Function**

Binds to scaffold/matrix attachment region (S/MAR) DNA. Can function as an estrogen receptor corepressor and can also inhibit cell proliferation.

Cellular Location

Cytoplasm. Nucleus.

Tissue Location

Expressed at high levels in the CNS and at low levels in the liver. Expressed in a wide number of breast cancer cell lines

SAFB2 Antibody (internal region) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

SAFB2 Antibody (internal region) - Images**SAFB2 Antibody (internal region) - Background**

This antibody is not expected to cross-react with SAFB1.

SAFB2 Antibody (internal region) - References

Scaffold attachment factors SAFB1 and SAFB2: Innocent bystanders or critical players in breast tumorigenesis?
players in breast tumorigenesis Oesterreich S. J Cell Biochem. 2003 Nov 1;90(4):653-61. PMID: 14587024