

MDFI Antibody (internal region)
Peptide-affinity purified goat antibody
Catalog # AF2976a**Specification**

MDFI Antibody (internal region) - Product Information

Application	E
Primary Accession	Q99750
Other Accession	NP_005577.1 , 4188
Predicted	Human
Host	Goat
Clonality	Polyclonal
Concentration	0.5 mg/ml
Isotype	IgG
Calculated MW	25029

MDFI Antibody (internal region) - Additional Information**Gene ID** 4188**Other Names**

MyoD family inhibitor, Myogenic repressor I-mf, MDFI

Dilution

E~~N/A

Format

0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin

Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

MDFI Antibody (internal region) is for research use only and not for use in diagnostic or therapeutic procedures.

MDFI Antibody (internal region) - Protein Information**Name** MDFI**Function**

Inhibits the transactivation activity of the Myod family of myogenic factors and represses myogenesis (By similarity). Acts by associating with Myod family members and retaining them in the cytoplasm by masking their nuclear localization signals (By similarity). Can also interfere with the DNA-binding activity of Myod family members (By similarity). Plays an important role in trophoblast and chondrogenic differentiation (By similarity). Regulates the transcriptional activity

of TCF7L1/TCF3 by interacting directly with TCF7L1/TCF3 and preventing it from binding DNA (By similarity). Binds to the axin complex, resulting in an increase in the level of free beta-catenin (By similarity). Affects axin regulation of the WNT and JNK signaling pathways (By similarity). Regulates the activity of mechanosensitive Piezo channel (PubMed:37590348).

Cellular Location

Nucleus. Cytoplasm {ECO:0000250|UniProtKB:P70331}

MDFI Antibody (internal region) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

MDFI Antibody (internal region) - Images**MDFI Antibody (internal region) - References**

Inhibitor of myogenic family, a novel suppressor of store-operated currents through an interaction with TRPC1. Ma R, Rundle D, Jacks J, Koch M, Downs T, Tsiokas L, The Journal of biological chemistry 2003 Dec 278 (52): 52763-72. PMID: 14530267