

**ICOS Antibody (internal region)**  
**Peptide-affinity purified goat antibody**  
**Catalog # AF2982a****Specification**

---

**ICOS Antibody (internal region) - Product Information**

Application	E
Primary Accession	<a href="#">O9Y6W8</a>
Other Accession	<a href="#">NP_036224.1</a> , <a href="#">29851</a> , <a href="#">54167 (mouse)</a>
Predicted	Human, Mouse
Host	Goat
Clonality	Polyclonal
Concentration	0.5 mg/ml
Isotype	IgG
Calculated MW	22625

**ICOS Antibody (internal region) - Additional Information****Gene ID** 29851**Other Names**

Inducible T-cell costimulator, Activation-inducible lymphocyte immunomediatory molecule, CD278, ICOS, AILIM

**Format**

0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin

**Storage**

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

**Precautions**

ICOS Antibody (internal region) is for research use only and not for use in diagnostic or therapeutic procedures.

**ICOS Antibody (internal region) - Protein Information****Name** ICOS**Synonyms** AILIM**Function**

Enhances all basic T-cell responses to a foreign antigen, namely proliferation, secretion of lymphokines, up-regulation of molecules that mediate cell-cell interaction, and effective help for antibody secretion by B-cells. Essential both for efficient interaction between T and B-cells and for normal antibody responses to T-cell dependent antigens. Does not up-regulate the production of interleukin- 2, but superinduces the synthesis of interleukin-10. Prevents the apoptosis of

pre-activated T-cells. Plays a critical role in CD40- mediated class switching of immunoglobulin isotypes (By similarity).

**Cellular Location**

[Isoform 1]: Cell membrane; Single- pass type I membrane protein

**Tissue Location**

Activated T-cells. Highly expressed on tonsillar T- cells, which are closely associated with B-cells in the apical light zone of germinal centers, the site of terminal B-cell maturation Expressed at lower levels in thymus, lung, lymph node and peripheral blood leukocytes. Expressed in the medulla of fetal and newborn thymus

**ICOS Antibody (internal region) - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

**ICOS Antibody (internal region) - Images****ICOS Antibody (internal region) - References**

Expression of ICOS on human melanoma-infiltrating CD4+CD25highFoxp3+ T regulatory cells: implications and impact on tumor-mediated immune suppression. Strauss L, Bergmann C, Szczepanski MJ, Lang S, Kirkwood JM, Whiteside TL, J Immunol. 2008 Mar 1;180(5):2967-80. PMID: 18292519