

NFATC4 / NFAT3 Antibody (internal region)
Peptide-affinity purified goat antibody
Catalog # AF2987a**Specification**

NFATC4 / NFAT3 Antibody (internal region) - Product Information

Application	E
Primary Accession	O14934
Other Accession	NP_001129494.1 , NP_004545.2 , 4776
Predicted	Human, Dog
Host	Goat
Clonality	Polyclonal
Concentration	0.5 mg/ml
Isotype	IgG
Calculated MW	95449

NFATC4 / NFAT3 Antibody (internal region) - Additional Information

Gene ID 4776

Other Names

Nuclear factor of activated T-cells, cytoplasmic 4, NF-ATc4, NFATc4, T-cell transcription factor NFAT3, NF-AT3, NFATC4, NFAT3

Format

0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin

Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

NFATC4 / NFAT3 Antibody (internal region) is for research use only and not for use in diagnostic or therapeutic procedures.

NFATC4 / NFAT3 Antibody (internal region) - Protein Information

Name NFATC4

Function

Ca(2+)-regulated transcription factor that is involved in several processes, including the development and function of the immune, cardiovascular, musculoskeletal, and nervous systems (PubMed: [11514544](http://www.uniprot.org/citations/11514544)), PubMed: [11997522](http://www.uniprot.org/citations/11997522), PubMed: [17213202](http://www.uniprot.org/citations/17213202), PubMed: [17875713](http://www.uniprot.org/citations/17875713), PubMed: [18668201](http://www.uniprot.org/citations/18668201)),

PubMed: 25663301, PubMed: 7749981). Involved in T-cell activation, stimulating the transcription of cytokine genes, including that of IL2 and IL4 (PubMed: 18347059, PubMed: 18668201, PubMed: 7749981). Along with NFATC3, involved in embryonic heart development. Following JAK/STAT signaling activation and as part of a complex with NFATC3 and STAT3, binds to the alpha-beta E4 promoter region of CRYAB and activates transcription in cardiomyocytes (By similarity). Involved in mitochondrial energy metabolism required for cardiac morphogenesis and function (By similarity). Transactivates many genes involved in the cardiovascular system, including AGTR2, NPPB/BNP (in synergy with GATA4), NPPA/ANP/ANF and MYH7/beta-MHC (By similarity). Involved in the regulation of adult hippocampal neurogenesis. Involved in BDNF-driven pro-survival signaling in hippocampal adult-born neurons. Involved in the formation of long-term spatial memory and long-term potentiation (By similarity). In cochlear nucleus neurons, may play a role in deafferentation-induced apoptosis during the developmental critical period, when auditory neurons depend on afferent input for survival (By similarity). Binds to and activates the BACE1/Beta-secretase 1 promoter, hence may regulate the proteolytic processing of the amyloid precursor protein (APP) (PubMed: 25663301). Plays a role in adipocyte differentiation (PubMed: 11997522). May be involved in myoblast differentiation into myotubes (PubMed: 17213202). Binds the consensus DNA sequence 5'-GGAAAAT-3' (Probable). In the presence of CREBBP, activates TNF transcription (PubMed: 11514544). Binds to PPARG gene promoter and regulates its activity (PubMed: 11997522). Binds to PPARG and REG3G gene promoters (By similarity).

Cellular Location

Cytoplasm. Nucleus. Note=When hyperphosphorylated, localizes in the cytosol. When intracellular Ca(2+) levels increase, dephosphorylation by calcineurin/PPP3CA leads to translocation into the nucleus (PubMed:11997522, PubMed:18347059). MAPK7/ERK5 and MTOR regulate NFATC4 nuclear export through phosphorylation at Ser-168 and Ser-170 (PubMed:18347059).

Tissue Location

Widely expressed, with high levels in placenta, lung, kidney, testis and ovary (PubMed:18675896). Weakly expressed in spleen and thymus (PubMed:18675896). In the hippocampus, expressed in the granular layer of the dentate gyrus, in the pyramidal neurons of CA3 region, and in the hippocampal fissure (PubMed:18675896). Expressed in the heart (at protein level) (PubMed:12370307)

NFATC4 / NFAT3 Antibody (internal region) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

NFATC4 / NFAT3 Antibody (internal region) - Images

NFATC4 / NFAT3 Antibody (internal region) - Background

This antibody is expected to recognize both isoforms (NP_001129494.1; NP_004545.2).

NFATC4 / NFAT3 Antibody (internal region) - References

Nuclear factor of activated T3 is a negative regulator of Ras-JNK1/2-AP-1 induced cell transformation. Yao K, Cho YY, Bergen HR, Madden BJ, Choi BY, Ma WY, Bode AM, Dong Z, Cancer Res. 2007 Sep 67 (18): 8725-35. PMID: 17875713