

eIF5A Antibody (internal region, near N-Term)
Peptide-affinity purified goat antibody
Catalog # AF3005a**Specification**

eIF5A Antibody (internal region, near N-Term) - Product Information

Application	WB, IHC, Pep-ELISA
Primary Accession	P63241
Other Accession	NP_001137232.1 , 1984
Reactivity	Human
Host	Goat
Clonality	Polyclonal
Concentration	0.5 mg/ml
Isotype	IgG
Calculated MW	16832

eIF5A Antibody (internal region, near N-Term) - Additional Information**Gene ID** 1984**Other Names**

Eukaryotic translation initiation factor 5A-1, eIF-5A-1, eIF-5A1, Eukaryotic initiation factor 5A isoform 1, eIF-5A, Rev-binding factor, eIF-4D, EIF5A

Dilution

WB~~1:1000
IHC~~1:100~500
Pep-ELISA~~N/A

Format

0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin

Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

eIF5A Antibody (internal region, near N-Term) is for research use only and not for use in diagnostic or therapeutic procedures.

eIF5A Antibody (internal region, near N-Term) - Protein Information**Name** EIF5A ([HGNC:3300](#))**Function**

Translation factor that promotes translation elongation and termination, particularly upon ribosome stalling at specific amino acid sequence contexts (PubMed:<a

[33547280](http://www.uniprot.org/citations/33547280)). Binds between the exit (E) and peptidyl (P) site of the ribosome and promotes rescue of stalled ribosome: specifically required for efficient translation of polyproline-containing peptides as well as other motifs that stall the ribosome (By similarity). Acts as a ribosome quality control (RQC) cofactor by joining the RQC complex to facilitate peptidyl transfer during CAT tailing step (By similarity). Also involved in actin dynamics and cell cycle progression, mRNA decay and probably in a pathway involved in stress response and maintenance of cell wall integrity (PubMed:[16987817](http://www.uniprot.org/citations/16987817)). With syntenin SDCBP, functions as a regulator of p53/TP53 and p53/TP53-dependent apoptosis (PubMed:[15371445](http://www.uniprot.org/citations/15371445)). Also regulates TNF-alpha-mediated apoptosis (PubMed:[15452064](http://www.uniprot.org/citations/15452064), PubMed:[17187778](http://www.uniprot.org/citations/17187778)). Mediates effects of polyamines on neuronal process extension and survival (PubMed:[17360499](http://www.uniprot.org/citations/17360499)). Is required for autophagy by assisting the ribosome in translating the ATG3 protein at a specific amino acid sequence, the 'ASP-ASP-Gly' motif, leading to the increase of the efficiency of ATG3 translation and facilitation of LC3B lipidation and autophagosome formation (PubMed:[29712776](http://www.uniprot.org/citations/29712776)).

Cellular Location

Cytoplasm. Nucleus. Endoplasmic reticulum membrane; Peripheral membrane protein; Cytoplasmic side. Note=Hypusine modification promotes the nuclear export and cytoplasmic localization and there was a dynamic shift in the localization from predominantly cytoplasmic to primarily nuclear under apoptotic inducing conditions (PubMed:19379712, PubMed:27306458). Nuclear export of hypusinated protein is mediated by XPO4 (PubMed:10944119, PubMed:27306458).

Tissue Location


Expressed in umbilical vein endothelial cells and several cancer cell lines (at protein level)

eIF5A Antibody (internal region, near N-Term) - Protocols

Provided below are standard protocols that you may find useful for product applications.


- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

eIF5A Antibody (internal region, near N-Term) - Images







EB09590 (0.1 μ g/ml) staining of MOLT4 lysate (35 μ g protein in RIPA buffer). Detected by chemiluminescence.



EB09590 (3.75 μ g/ml) staining of paraffin embedded Human Spleen. Steamed antigen retrieval with citrate buffer pH 6, AP-staining.



EB09590 (3.75 μ g/ml) staining of paraffin embedded Human Tonsil. Steamed antigen retrieval with citrate buffer pH 6, AP-staining.



EB09590 (3.75µg/ml) staining of paraffin embedded Human Vessel. Steamed antigen retrieval with citrate buffer pH 6, AP-staining.

eIF5A Antibody (internal region, near N-Term) - Background

Reported variants NP_001137233.1, NP_001961.1 and NP_001137234.1 represent identical protein:

eIF5A Antibody (internal region, near N-Term) - References

The genomic structure encoding human initiation factor eIF-5A Koettwitz K, Kappel B, Baumruker T, Hauber J, Bevec D Gene. 1994 Jul 144 (2): 249-52 PMID: 7545941