

**WASF1 / WAVE Antibody (internal region)**  
**Peptide-affinity purified goat antibody**  
**Catalog # AF3075a****Specification**

---

**WASF1 / WAVE Antibody (internal region) - Product Information**

Application	E
Primary Accession	<a href="#">O92558</a>
Other Accession	<a href="#">NP_003922.1</a> , <a href="#">8936</a> , <a href="#">83767 (mouse)</a> , <a href="#">294568 (rat)</a>
Predicted	Human, Mouse, Rat, Dog
Host	Goat
Clonality	Polyclonal
Concentration	0.5 mg/ml
Isotype	IgG
Calculated MW	61652

**WASF1 / WAVE Antibody (internal region) - Additional Information****Gene ID** 8936**Other Names**

Wiskott-Aldrich syndrome protein family member 1, WASP family protein member 1, Protein WAVE-1, Verprolin homology domain-containing protein 1, WASF1, KIAA0269, SCAR1, WAVE1

**Dilution**

E~~N/A

**Format**

0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin

**Storage**

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

**Precautions**

WASF1 / WAVE Antibody (internal region) is for research use only and not for use in diagnostic or therapeutic procedures.

**WASF1 / WAVE Antibody (internal region) - Protein Information****Name** WASF1 ([HGNC:12732](#))**Synonyms** KIAA0269, SCAR1, WAVE1**Function**

Downstream effector molecule involved in the transmission of signals from tyrosine kinase

receptors and small GTPases to the actin cytoskeleton. Promotes formation of actin filaments. Part of the WAVE complex that regulates lamellipodia formation (PubMed:<a href="http://www.uniprot.org/citations/29961568" target="\_blank">29961568</a>). The WAVE complex regulates actin filament reorganization via its interaction with the Arp2/3 complex (By similarity). As component of the WAVE1 complex, required for BDNF-NTRK2 endocytic trafficking and signaling from early endosomes (By similarity). Also involved in the regulation of mitochondrial dynamics (PubMed:<a href="http://www.uniprot.org/citations/29961568" target="\_blank">29961568</a>).

#### **Cellular Location**

Cytoplasm, cytoskeleton. Synapse {ECO:0000250|UniProtKB:Q5BJU7} Cell junction, focal adhesion. Note=Dot- like pattern in the cytoplasm. Concentrated in Rac-regulated membrane-ruffling areas (PubMed:9889097). Partial translocation to focal adhesion sites might be mediated by interaction with SORBS2 (PubMed:18559503). In neurons, colocalizes with activated NTRK2 after BDNF addition in endocytic sites through the association with TMEM108 (By similarity). {ECO:0000250|UniProtKB:Q8R5H6, ECO:0000269|PubMed:18559503, ECO:0000269|PubMed:9889097}

#### **Tissue Location**

Highly expressed in brain. Lowly expressed in testis, ovary, colon, kidney, pancreas, thymus, small intestine and peripheral blood

### **WASF1 / WAVE Antibody (internal region) - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

### **WASF1 / WAVE Antibody (internal region) - Images**

### **WASF1 / WAVE Antibody (internal region) - Background**

Reported variants represent identical protein: NP\_001020106.1, NP\_003922.1, NP\_001020107.1, NP\_001020105.1

### **WASF1 / WAVE Antibody (internal region) - References**

WAVE and Arp2/3 jointly inhibit filopodium formation by entering into a complex with mDia2. Beli P, Mascheroni D, Xu D, Innocenti M. Nature cell biology 2008 Jul 10 (7): 849-57. PMID: 18516090