

POU2F2 / OCT2 Antibody (N-Term)
Peptide-affinity purified goat antibody
Catalog # AF3196a**Specification**

POU2F2 / OCT2 Antibody (N-Term) - Product Information

Application	WB, E
Primary Accession	P09086
Other Accession	NP_002689.1 , 5452
Reactivity	Human, Mouse
Predicted	Rat, Pig, Dog
Host	Goat
Clonality	Polyclonal
Concentration	0.5 mg/ml
Isotype	IgG
Calculated MW	51209

POU2F2 / OCT2 Antibody (N-Term) - Additional Information**Gene ID** 5452**Other Names**

POU domain, class 2, transcription factor 2, Lymphoid-restricted immunoglobulin octamer-binding protein NF-A2, Octamer-binding protein 2, Oct-2, Octamer-binding transcription factor 2, OTF-2, POU2F2, OCT2, OTF2

Dilution

WB~~1:1000

E~~N/A

Format

0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin

Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

POU2F2 / OCT2 Antibody (N-Term) is for research use only and not for use in diagnostic or therapeutic procedures.

POU2F2 / OCT2 Antibody (N-Term) - Protein Information**Name** POU2F2 ([HGNC:9213](#))**Synonyms** OCT2, OTF2

Function

Transcription factor that specifically binds to the octamer motif (5'-ATTTGCAT-3') (PubMed:2904654, PubMed:7859290). Regulates IL6 expression in B cells with POU2AF1 (By similarity). Regulates transcription in a number of tissues in addition to activating immunoglobulin gene expression (PubMed:2901913, PubMed:2904654). Modulates transcription transactivation by NR3C1, AR and PGR (PubMed:10480874).

Cellular Location

Cytoplasm. Nucleus.

Tissue Location

Isoform 3 is B-cell specific. Isoform 5 is expressed in B-cells and the immunoglobulin-expressing T-cell line MOLT-4, but not in the T-cell line BW5147.

POU2F2 / OCT2 Antibody (N-Term) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

POU2F2 / OCT2 Antibody (N-Term) - Images

AF3196a (0.1 µg/ml) staining of Human Tonsil lysate (35 µg protein in RIPA buffer). Primary incubation was 1 hour. Detected by chemiluminescence.

POU2F2 / OCT2 Antibody (N-Term) - References

Functional long-range interactions of the IgH 3' enhancers with the bcl-2 promoter region in t(14;18) lymphoma cells. Duan H, Xiang H, Ma L, Boxer LM, Oncogene 2008 Dec 27 (53): 6720-8. PMID: 18695675