

NOX1 Antibody (internal region)
Peptide-affinity purified goat antibody
Catalog # AF3415a**Specification**

NOX1 Antibody (internal region) - Product Information

Application	WB, E
Primary Accession	Q9Y5S8
Other Accession	NP_008983.2 , NP_39249.1 , 27035 , 237038 (mouse)
Reactivity	Human, Mouse
Predicted	Dog
Host	Goat
Clonality	Polyclonal
Concentration	0.5 mg/ml
Isotype	IgG
Calculated MW	64871

NOX1 Antibody (internal region) - Additional Information**Gene ID** 27035**Other Names**

NADPH oxidase 1, NOX-1, 1.-.-., Mitogenic oxidase 1, MOX-1, NADH/NADPH mitogenic oxidase subunit P65-MOX, NOH-1, NOX1, MOX1, NOH1

Dilution

WB~~1:1000

E~~N/A

Format

0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin

Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

NOX1 Antibody (internal region) is for research use only and not for use in diagnostic or therapeutic procedures.

NOX1 Antibody (internal region) - Protein Information**Name** NOX1 ([HGNC:7889](#))**Synonyms** MOX1, NOH1

Function

NADPH oxidase that catalyzes the generation of superoxide from molecular oxygen utilizing NADPH as an electron donor.

Cellular Location

Cell projection, invadopodium membrane; Multi-pass membrane protein. Cell membrane; Multi-pass membrane protein


Tissue Location

[Isoform NOH-1L]: Detected in colon, uterus, prostate, and colon carcinoma, but not in peripheral blood leukocytes

NOX1 Antibody (internal region) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

NOX1 Antibody (internal region) - Images

AF3415a (0.1µg/ml) staining of lysate of cell line colorectal adenocarcinoma (A) and HeLa (B) (35µg protein in RIPA buffer). Primary incubation was 1 hour. Detected by chemiluminescence.

NOX1 Antibody (internal region) - Background

This antibody is expected to recognize both reported isoforms (NP_008983.2; NP_39249.1).

NOX1 Antibody (internal region) - References

Transcriptional regulation of NADPH oxidase isoforms, Nox1 and Nox4, by nuclear factor-kappaB in

human aortic smooth muscle cells. Manea A, Tanase LI, Raicu M, Simionescu M, Biochemical and biophysical research communications 2010 Jun 396 (4): 901-7. PMID: 20457132