

Mucin 10 / Prol1 Antibody (internal region)
Peptide-affinity purified goat antibody
Catalog # AF3427a**Specification**

Mucin 10 / Prol1 Antibody (internal region) - Product Information

Application	E
Primary Accession	E9PYQ4
Other Accession	NP_032670.2 , 17830 (mouse)
Predicted	Mouse
Host	Goat
Clonality	Polyclonal
Concentration	0.5 mg/ml
Isotype	IgG

Mucin 10 / Prol1 Antibody (internal region) - Additional Information**Other Names**

Prol1; proline rich, lacrimal 1; Muc10; mucin 10; mucin 10, submandibular gland salivary mucin

Format

0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin

Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

Mucin 10 / Prol1 Antibody (internal region) is for research use only and not for use in diagnostic or therapeutic procedures.

Mucin 10 / Prol1 Antibody (internal region) - Protein Information**Mucin 10 / Prol1 Antibody (internal region) - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Mucin 10 / Prol1 Antibody (internal region) - Images

Mucin 10 / Prol1 Antibody (internal region) - References

Human Opiorphin, a natural antinociceptive modulator of opioid-dependent pathways. Wisner A, Dufour E, Messaoudi M, Nejdi A, Marcel A, Ungeheuer MN, Rougeot C, Proceedings of the National Academy of Sciences of the United States of America 2006 Nov 103 (47): 17979-84. PMID: 17101991