

Mucin 19 / Apomucin Antibody (C-Term)
Peptide-affinity purified goat antibody
Catalog # AF3430a**Specification**

Mucin 19 / Apomucin Antibody (C-Term) - Product Information

Application	IHC, WB
Primary Accession	Q6PZE0.2
Other Accession	NP_997126.2 , 239611 (mouse) , 497227 (rat)
Reactivity	Mouse
Predicted	Rat
Host	Goat
Clonality	Polyclonal
Concentration	0.5 mg/ml
Isotype	IgG

Mucin 19 / Apomucin Antibody (C-Term) - Additional Information**Other Names**

Muc19; mucin 19; MUC-19; gel-forming secreted mucin-19; mucin-19; sublingual apomucin; mucin apoprotein

Format

0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin

Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

Mucin 19 / Apomucin Antibody (C-Term) is for research use only and not for use in diagnostic or therapeutic procedures.


Mucin 19 / Apomucin Antibody (C-Term) - Protein Information**Mucin 19 / Apomucin Antibody (C-Term) - Protocols**

Provided below are standard protocols that you may find useful for product applications.


- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)

- [Cell Culture](#)

Mucin 19 / Apomucin Antibody (C-Term) - Images



EB10619 (1.8ug/ml) staining cells of the sublingual salivary gland (SLG) in mouse, but not any cells of the submandibular salivary gland (SMG).



EB10619 (0.25 µg/ml) staining of homogenized Mouse salivary gland tissue. (20 µg protein per lane). Detected by chemiluminescence.

Mucin 19 / Apomucin Antibody (C-Term) - References

Detailed haplotype-tagging study of germline variation of MUC19 in inflammatory bowel disease. Phillips AM, Nimmo ER, Van Limbergen J, Drummond HE, Smith L, Satsangi J, Inflammatory bowel diseases 2010 Apr 16 (4): 557-8. PMID: 19714762