

Cd28 (mouse) Antibody (internal region)
Peptide-affinity purified goat antibody
Catalog # AF3470a**Specification**

Cd28 (mouse) Antibody (internal region) - Product Information

Application	WB
Primary Accession	P31041.2
Other Accession	NP_031668.3 , 12487 (mouse)
Reactivity	Mouse
Host	Goat
Clonality	Polyclonal
Concentration	0.5 mg/ml
Isotype	IgG

Cd28 (mouse) Antibody (internal region) - Additional Information**Other Names**

Cd28; CD28 antigen; CH29-189K17.3; OTTMUSP00000002157; T-cell-specific surface glycoprotein CD28

Format

0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin

Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.


Precautions

Cd28 (mouse) Antibody (internal region) is for research use only and not for use in diagnostic or therapeutic procedures.

Cd28 (mouse) Antibody (internal region) - Protein Information**Cd28 (mouse) Antibody (internal region) - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Cd28 (mouse) Antibody (internal region) - Images

AF3470a (0.3 µg/ml) staining of Mouse Thymus lysate (35 µg protein in RIPA buffer). Primary incubation was 1 hour. Detected by chemiluminescence.

Cd28 (mouse) Antibody (internal region) - Background

The immunizing peptide represents a part of the extracellular domain.

Cd28 (mouse) Antibody (internal region) - References

New approach reveals CD28 and IFNG gene interaction in the susceptibility to cervical cancer.
Guzman VB, Yambartsev A, Goncalves-Primo A, Silva ID, Carvalho CR, Ribalta JC, Goulart LR, Shulzhenko N, Gerbase-Delima M, Morgun A, Human molecular genetics 2008 Jun 17 (12): 1838-44. PMID: 18337305