

Ghrelin O-acyltransferase Antibody (internal region)

Peptide-affinity purified goat antibody Catalog # AF3497a

Specification

Ghrelin O-acyltransferase Antibody (internal region) - Product Information

Application

Primary Accession <u>Q96T53</u>

Other Accession <u>NP_001094386.1</u>, <u>619373</u>

Predicted Human
Host Goat
Clonality Polyclonal
Concentration 0.5 mg/ml

Isotype IgG
Calculated MW 49716

Ghrelin O-acyltransferase Antibody (internal region) - Additional Information

Gene ID 619373

Other Names

Ghrelin O-acyltransferase, 2.3.1.-, Membrane-bound O-acyltransferase domain-containing protein 4, O-acyltransferase domain-containing protein 4, MBOAT4, GOAT, OACT4

Dilution

E~~N/A

Format

0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin

Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

Ghrelin O-acyltransferase Antibody (internal region) is for research use only and not for use in diagnostic or therapeutic procedures.

Ghrelin O-acyltransferase Antibody (internal region) - Protein Information

Name MBOAT4 (HGNC:32311)

Synonyms GOAT, OACT4

Function

Catalyzes ghrelin acylation at 'Ser-3' using preferentially octanoyl-CoA, hexanoyl-CoA and decanoyl-CoA as acyl-CoA donors leading to ghrelin activity (PubMed:<a



Tel: 858.875.1900 Fax: 858.875.1999

href="http://www.uniprot.org/citations/18443287" target=" blank">18443287, PubMed:24045953, PubMed:25562443, PubMed:28134508). In vitro also uses acyl-CoA donors of different lengths from short-chain (C2) to long-chain fatty acids (C16) knowing that acyl-CoA donors from butanoyl-CoA (C4) to dodecanoyl-CoA (C12) are more efficient compared to longer acyl-CoA donors, such as myristoyl- CoA (C14) and palmitoyl-CoA (C16) that are not efficient (PubMed: 18443287).

Cellular Location

Endoplasmic reticulum membrane {ECO:0000250|UniProtKB:P0C7A3}; Multi-pass membrane protein {ECO:0000250|UniProtKB:P0C7A3}

Tissue Location

Expressed predominantly in stomach with moderate levels in pancreas and relatively low levels in most other tissues

Ghrelin O-acyltransferase Antibody (internal region) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- <u>Immunohistochemistry</u>
- <u>Immunofluorescence</u>
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

Ghrelin O-acyltransferase Antibody (internal region) - Images

Ghrelin O-acyltransferase Antibody (internal region) - References

Production of n-octanoyl-modified ghrelin in cultured cells requires prohormone processing protease and ghrelin O-acyltransferase, as well as n-octanoic acid. Takahashi T, Ida T, Sato T, Nakashima Y, Nakamura Y, Tsuji A, Kojima M, Journal of biochemistry 2009 Nov 146 (5): 675-82. PMID: 19628676