

CBARA1 (aa113-126) Antibody (internal region)
Peptide-affinity purified goat antibody
Catalog # AF3517a

Specification

CBARA1 (aa113-126) Antibody (internal region) - Product Information

Application	WB, E
Primary Accession	Q9BPX6
Other Accession	NP_006068.2 , NP_001182447.1 , 10367 , 216001 (mouse) , 365567 (rat)
Reactivity	Human
Predicted	Mouse, Rat, Pig, Dog
Host	Goat
Clonality	Polyclonal
Concentration	0.5 mg/ml
Isotype	IgG
Calculated MW	54351

CBARA1 (aa113-126) Antibody (internal region) - Additional Information

Gene ID 10367

Other Names

Calcium uptake protein 1, mitochondrial, Atopy-related autoantigen CALC, ara CALC, Calcium-binding atopy-related autoantigen 1, Hom s 4, MICU1, CALC, CBARA1

Dilution

WB~~1:1000

E~~N/A

Format

0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin

Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

CBARA1 (aa113-126) Antibody (internal region) is for research use only and not for use in diagnostic or therapeutic procedures.

CBARA1 (aa113-126) Antibody (internal region) - Protein Information

Name MICU1 {ECO:0000303|PubMed:20693986, ECO:0000312|HGNC:HGNC:1530}

Function

Calcium sensor of the mitochondrial calcium uniporter (MCU) channel, which senses calcium level

via its EF-hand domains (PubMed:20693986, PubMed:23101630, PubMed:23747253, PubMed:24313810, PubMed:24332854, PubMed:24503055, PubMed:24560927, PubMed:26341627, PubMed:26903221, PubMed:27099988, PubMed:28615291, PubMed:30454562, PubMed:30638448, PubMed:32494073, PubMed:32667285, PubMed:32762847, PubMed:32790952, PubMed:34463251, PubMed:36206740, PubMed:37036971, PubMed:37126688). MICU1 and MICU2 (or MICU3) form a disulfide-linked heterodimer that stimulates and inhibits MCU activity, depending on the concentration of calcium (PubMed:24560927, PubMed:26903221, PubMed:28615291, PubMed:32148862, PubMed:32494073, PubMed:32667285, PubMed:32762847, PubMed:32790952, PubMed:36206740, PubMed:37036971, PubMed:37126688). At low calcium levels, MICU1 occludes the pore of the MCU channel, preventing mitochondrial calcium uptake (PubMed:32494073, PubMed:32667285, PubMed:32762847, PubMed:37036971, PubMed:37126688). At higher calcium levels, calcium-binding to MICU1 and MICU2 (or MICU3) induces a conformational change that weakens MCU-MICU1 interactions and moves the MICU1-MICU2 heterodimer away from the pore, allowing calcium permeation through the MCU channel (PubMed:32494073, PubMed:32667285, PubMed:32762847). Also required to protect against manganese toxicity by preventing manganese uptake by MCU: mechanistically, manganese-binding to its EF-hand domains does not induce any conformational change, maintaining MCU pore occlusion (PubMed:30082385, PubMed:30403999). Also acts as a barrier for inhibitors of the MCU channel, such as ruthenium red or its derivative Ru360 (PubMed:37244260). Acts as a regulator of mitochondrial cristae structure independently of its ability to regulate the mitochondrial calcium uniporter channel (PubMed:31427612, PubMed:37098122). Regulates

glucose-dependent insulin secretion in pancreatic beta-cells by regulating mitochondrial calcium uptake (PubMed:22904319). Induces T- helper 1-mediated autoreactivity, which is accompanied by the release of IFNG (PubMed:16002733).

Cellular Location

Mitochondrion intermembrane space. Mitochondrion inner membrane. Note=Recruited to the mitochondrial inner membrane by EMRE/SMDT1 (PubMed:30454562). Also localizes to mitochondrial cristae junctions (PubMed:31427612)

Tissue Location

Expressed in epithelial cell lines. Strongly expressed in epidermal keratinocytes and dermal endothelial cells

CBARA1 (aa113-126) Antibody (internal region) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

CBARA1 (aa113-126) Antibody (internal region) - Images



AF3517a (0.5 µg/ml) staining of K562 lysate (35 µg protein in RIPA buffer). Primary incubation was 1 hour. Detected by chemiluminescence.

CBARA1 (aa113-126) Antibody (internal region) - Background

This antibody is expected to recognize reported isoforms 1 and 2 (NP_006068.2; NP_001182447.1).

CBARA1 (aa113-126) Antibody (internal region) - References

MICU1 encodes a mitochondrial EF hand protein required for Ca(2+) uptake. Perocchi F, Gohil VM, Girgis HS, Bao XR, McCombs JE, Palmer AE, Mootha VK, Nature 2010 Sep 4; 467 (7313): 291-6. PMID: 20693986