

DNAJB9 (aa61-75) Antibody (internal region)
Peptide-affinity purified goat antibody
Catalog # AF3582a**Specification**

DNAJB9 (aa61-75) Antibody (internal region) - Product Information

Application	WB, E
Primary Accession	Q9UBS3
Other Accession	NP_036460.1 , 4189
Reactivity	Human, Mouse, Rat
Predicted	Pig, Dog
Host	Goat
Clonality	Polyclonal
Concentration	0.5 mg/ml
Isotype	IgG
Calculated MW	25518

DNAJB9 (aa61-75) Antibody (internal region) - Additional Information**Gene ID** 4189**Other Names**

DnaJ homolog subfamily B member 9, Endoplasmic reticulum DNA J domain-containing protein 4, ER-resident protein ERdj4, ERdj4, Microvascular endothelial differentiation gene 1 protein, Mdg-1, DNAJB9, MDG1

Dilution

WB~~1:1000

E~~N/A

Format

0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin

Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

DNAJB9 (aa61-75) Antibody (internal region) is for research use only and not for use in diagnostic or therapeutic procedures.

DNAJB9 (aa61-75) Antibody (internal region) - Protein Information**Name** DNAJB9**Synonyms** MDG1 {ECO:0000303|Ref.1}

Function

Co-chaperone for Hsp70 protein HSPA5/BiP that acts as a key repressor of the ERN1/IRE1-mediated unfolded protein response (UPR) (By similarity). J domain-containing co-chaperones stimulate the ATPase activity of Hsp70 proteins and are required for efficient substrate recognition by Hsp70 proteins (PubMed:18400946). In the unstressed endoplasmic reticulum, interacts with the luminal region of ERN1/IRE1 and selectively recruits HSPA5/BiP: HSPA5/BiP disrupts the dimerization of the active ERN1/IRE1 luminal region, thereby inactivating ERN1/IRE1 (By similarity). Also involved in endoplasmic reticulum-associated degradation (ERAD) of misfolded proteins. Required for survival of B- cell progenitors and normal antibody production (By similarity).

Cellular Location

Endoplasmic reticulum lumen {ECO:0000250|UniProtKB:Q9QYI6}


Tissue Location

Widely expressed. Expressed at highest level in the liver, placenta and kidney (PubMed:11836248)

DNAJB9 (aa61-75) Antibody (internal regoin) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

DNAJB9 (aa61-75) Antibody (internal regoin) - Images

AF3582a (1 µg/ml) staining of Human Liver lysate (35 µg protein in RIPA buffer). Primary incubation was 1 hour. Detected by chemiluminescence.

DNAJB9 (aa61-75) Antibody (internal regoin) - References

Assignment of the microvascular endothelial differentiation gene 1 (MDG1) to human chromosome band 14q24.2-->q24.3 by fluorescence in situ hybridization. Pröls F, Liehr T, Rinke R, Rautenstrauss B. Cytogenet Cell Genet. 1997;79(1-2):149-50. PMID: 9533036