

Histamine Receptor H2 (mouse) Antibody (internal region)
Peptide-affinity purified goat antibody
Catalog # AF3586a

Specification

Histamine Receptor H2 (mouse) Antibody (internal region) - Product Information

Application	E
Primary Accession	P97292.2
Other Accession	NP_001010973.1 , NP_032312.2 , 15466 (mouse) , 25461 (rat)
Predicted	Mouse, Rat
Host	Goat
Clonality	Polyclonal
Concentration	0.5 mg/ml
Isotype	IgG

Histamine Receptor H2 (mouse) Antibody (internal region) - Additional Information

Other Names

Hrh2; histamine receptor H2; H2R; HH2R; gastric receptor 1; gastric receptor I; histamine H2 receptor

Format

0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin

Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

Histamine Receptor H2 (mouse) Antibody (internal region) is for research use only and not for use in diagnostic or therapeutic procedures.

Histamine Receptor H2 (mouse) Antibody (internal region) - Protein Information

Histamine Receptor H2 (mouse) Antibody (internal region) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)

- [Cell Culture](#)

Histamine Receptor H2 (mouse) Antibody (internal region) - Images**Histamine Receptor H2 (mouse) Antibody (internal region) - Background**

This antibody is expected to recognize both reported isoforms (NP_001124527.1; NP_071640.1).

Histamine Receptor H2 (mouse) Antibody (internal region) - References

Histamine: its novel role as an endogenous regulator of Con A-dependent T cell proliferation.
Nakane H, Sonobe Y, Watanabe T, Nakano K. Inflamm Res. 2004 Jul;53(7):324-8. PMID: 15241568