

**Wfdc1 (mouse) Antibody (internal region , near N-Term)**  
**Peptide-affinity purified goat antibody**  
**Catalog # AF3588a****Specification**

---

**Wfdc1 (mouse) Antibody (internal region , near N-Term) - Product Information**

|                   |                                                                                            |
|-------------------|--------------------------------------------------------------------------------------------|
| Application       | WB                                                                                         |
| Primary Accession | <a href="#">Q9ESH5.1</a>                                                                   |
| Other Accession   | <a href="#">NP_075884.1</a> , <a href="#">67866 (mouse)</a> , <a href="#">171112 (rat)</a> |
| Reactivity        | Mouse                                                                                      |
| Predicted         | Rat                                                                                        |
| Host              | Goat                                                                                       |
| Clonality         | Polyclonal                                                                                 |
| Concentration     | 0.5 mg/ml                                                                                  |
| Isotype           | IgG                                                                                        |

**Wfdc1 (mouse) Antibody (internal region , near N-Term) - Additional Information****Other Names**

WFDC1; WAP four-disulfide core domain 1; ps20; WAP four-disulfide core domain 1 homolog; WAP four-disulfide core domain protein 1; prostate stromal protein ps20; ps20 growth inhibitor; 2310058A03Rik

**Format**

0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin

**Storage**

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

**Precautions**

Wfdc1 (mouse) Antibody (internal region , near N-Term) is for research use only and not for use in diagnostic or therapeutic procedures.

**Wfdc1 (mouse) Antibody (internal region , near N-Term) - Protein Information****Wfdc1 (mouse) Antibody (internal region , near N-Term) - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)

- [Flow Cytometry](#)
- [Cell Culture](#)

#### **Wfdc1 (mouse) Antibody (internal region , near N-Term) - Images**



AF3588a (0.3 µg/ml) staining of Mouse Heart lysate (35 µg protein in RIPA buffer). Primary incubation was 1 hour. Detected by chemiluminescence.

#### **Wfdc1 (mouse) Antibody (internal region , near N-Term) - References**

Molecular analysis of WFDC1/ps20 gene in prostate cancer. Watson JE, Kamkar S, James K, Kowbel D, Andaya A, Paris PL, Simko J, Carroll P, McAlhany S, Rowley D, Collins C. Prostate. 2004 Oct 1;61(2):192-9. PMID: 15305342