

**CHUK Antibody (internal region, near C-Term)**  
**Peptide-affinity purified goat antibody**  
**Catalog # AF3714a****Specification**

---

**CHUK Antibody (internal region, near C-Term) - Product Information**

Application	WB, E
Primary Accession	<a href="#">O15111</a>
Other Accession	<a href="#">NP_001269.3</a> , <a href="#">1147</a>
Reactivity	Human
Host	Goat
Clonality	Polyclonal
Concentration	0.5 mg/ml
Isotype	IgG
Calculated MW	84640

**CHUK Antibody (internal region, near C-Term) - Additional Information****Gene ID** 1147**Other Names**

Inhibitor of nuclear factor kappa-B kinase subunit alpha, I-kappa-B kinase alpha, IKK-A, IKK-alpha, IkbKA, IkappaB kinase, 2.7.11.10, Conserved helix-loop-helix ubiquitous kinase, I-kappa-B kinase 1, IKK1, Nuclear factor NF-kappa-B inhibitor kinase alpha, NFKBKA, Transcription factor 16, TCF-16, CHUK, IKKA, TCF16

**Dilution**

WB~~~1:1000

E~~N/A

**Format**

0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin

**Storage**

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

**Precautions**

CHUK Antibody (internal region, near C-Term) is for research use only and not for use in diagnostic or therapeutic procedures.

**CHUK Antibody (internal region, near C-Term) - Protein Information****Name** CHUK**Synonyms** IKKA, TCF16

## Function

Serine kinase that plays an essential role in the NF-kappa-B signaling pathway which is activated by multiple stimuli such as inflammatory cytokines, bacterial or viral products, DNA damages or other cellular stresses (PubMed:<a href="http://www.uniprot.org/citations/18626576" target="\_blank">18626576</a>, PubMed:<a href="http://www.uniprot.org/citations/9244310" target="\_blank">9244310</a>, PubMed:<a href="http://www.uniprot.org/citations/9252186" target="\_blank">9252186</a>, PubMed:<a href="http://www.uniprot.org/citations/9346484" target="\_blank">9346484</a>). Acts as a part of the canonical IKK complex in the conventional pathway of NF-kappa-B activation and phosphorylates inhibitors of NF-kappa-B on serine residues (PubMed:<a href="http://www.uniprot.org/citations/18626576" target="\_blank">18626576</a>, PubMed:<a href="http://www.uniprot.org/citations/35952808" target="\_blank">35952808</a>, PubMed:<a href="http://www.uniprot.org/citations/9244310" target="\_blank">9244310</a>, PubMed:<a href="http://www.uniprot.org/citations/9252186" target="\_blank">9252186</a>, PubMed:<a href="http://www.uniprot.org/citations/9346484" target="\_blank">9346484</a>). These modifications allow polyubiquitination of the inhibitors and subsequent degradation by the proteasome (PubMed:<a href="http://www.uniprot.org/citations/18626576" target="\_blank">18626576</a>, PubMed:<a href="http://www.uniprot.org/citations/9244310" target="\_blank">9244310</a>, PubMed:<a href="http://www.uniprot.org/citations/9252186" target="\_blank">9252186</a>, PubMed:<a href="http://www.uniprot.org/citations/9346484" target="\_blank">9346484</a>). In turn, free NF-kappa-B is translocated into the nucleus and activates the transcription of hundreds of genes involved in immune response, growth control, or protection against apoptosis (PubMed:<a href="http://www.uniprot.org/citations/18626576" target="\_blank">18626576</a>, PubMed:<a href="http://www.uniprot.org/citations/9244310" target="\_blank">9244310</a>, PubMed:<a href="http://www.uniprot.org/citations/9252186" target="\_blank">9252186</a>, PubMed:<a href="http://www.uniprot.org/citations/9346484" target="\_blank">9346484</a>). Negatively regulates the pathway by phosphorylating the scaffold protein TAXBP1 and thus promoting the assembly of the A20/TNFAIP3 ubiquitin-editing complex (composed of A20/TNFAIP3, TAX1BP1, and the E3 ligases ITCH and RNF11) (PubMed:<a href="http://www.uniprot.org/citations/21765415" target="\_blank">21765415</a>). Therefore, CHUK plays a key role in the negative feedback of NF-kappa-B canonical signaling to limit inflammatory gene activation. As part of the non-canonical pathway of NF-kappa-B activation, the MAP3K14-activated CHUK/IKKA homodimer phosphorylates NFkB2/p100 associated with RelB, inducing its proteolytic processing to NFkB2/p52 and the formation of NF-kappa-B RelB-p52 complexes (PubMed:<a href="http://www.uniprot.org/citations/20501937" target="\_blank">20501937</a>). In turn, these complexes regulate genes encoding molecules involved in B-cell survival and lymphoid organogenesis. Also participates in the negative feedback of the non-canonical NF-kappa-B signaling pathway by phosphorylating and destabilizing MAP3K14/NIK. Within the nucleus, phosphorylates CREBBP and consequently increases both its transcriptional and histone acetyltransferase activities (PubMed:<a href="http://www.uniprot.org/citations/17434128" target="\_blank">17434128</a>). Modulates chromatin accessibility at NF-kappa-B- responsive promoters by phosphorylating histones H3 at 'Ser-10' that are subsequently acetylated at 'Lys-14' by CREBBP (PubMed:<a href="http://www.uniprot.org/citations/12789342" target="\_blank">12789342</a>). Additionally, phosphorylates the CREBBP-interacting protein NCOA3. Also phosphorylates FOXO3 and may regulate this pro-apoptotic transcription factor (PubMed:<a href="http://www.uniprot.org/citations/15084260" target="\_blank">15084260</a>). Phosphorylates RIPK1 at 'Ser-25' which represses its kinase activity and consequently prevents TNF-mediated RIPK1-dependent cell death (By similarity). Phosphorylates AMBRA1 following mitophagy induction, promoting AMBRA1 interaction with ATG8 family proteins and its mitophagic activity (PubMed:<a href="http://www.uniprot.org/citations/30217973" target="\_blank">30217973</a>).

## Cellular Location

Cytoplasm. Nucleus Note=Shuttles between the cytoplasm and the nucleus

## Tissue Location

Widely expressed.

**CHUK Antibody (internal region, near C-Term) - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

**CHUK Antibody (internal region, near C-Term) - Images**

AF3714a (1  $\mu$ g/ml) staining of HeLa nuclear lysate (35  $\mu$ g protein in RIPA buffer). Primary incubation was 1 hour. Detected by chemiluminescence.

**CHUK Antibody (internal region, near C-Term) - References**

How NF-kappaB is activated: the role of the IkappaB kinase (IKK) complex. Karin M. Oncogene. 1999 Nov 22;18(49):6867-74. PMID: 10602462