

NCKAP1 Antibody (internal region)
Peptide-affinity purified goat antibody
Catalog # AF3839a**Specification**

NCKAP1 Antibody (internal region) - Product Information

Application	E
Primary Accession	O9Y2A7
Other Accession	NP_038464.1 , NP_995314.1 , 10787 , 50884 (mouse) , 58823 (rat)
Predicted Host	Human, Mouse, Rat, Pig, Dog
Clonality	Goat
Concentration	Polyclonal
Isotype	0.5 mg/ml
Calculated MW	IgG
	128790

NCKAP1 Antibody (internal region) - Additional Information**Gene ID** 10787**Other Names**

Nck-associated protein 1, NAP 1, Membrane-associated protein HEM-2, p125Nap1, NCKAP1, HEM2, KIAA0587, NAP1

Dilution

E~~N/A

Format

0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin

Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

NCKAP1 Antibody (internal region) is for research use only and not for use in diagnostic or therapeutic procedures.

NCKAP1 Antibody (internal region) - Protein Information**Name** NCKAP1**Synonyms** HEM2, KIAA0587, NAP1**Function**

Part of the WAVE complex that regulates lamellipodia formation. The WAVE complex regulates

actin filament reorganization via its interaction with the Arp2/3 complex. Actin remodeling activity is regulated by RAC1. As component of the WAVE1 complex, required for BDNF-NTRK2 endocytic trafficking and signaling from early endosomes.

Cellular Location

Cell membrane {ECO:0000250|UniProtKB:P28660}; Single-pass membrane protein {ECO:0000250|UniProtKB:P28660}; Cytoplasmic side {ECO:0000250|UniProtKB:P28660}. Cell projection, lamellipodium membrane {ECO:0000250|UniProtKB:P28660}; Single-pass membrane protein {ECO:0000250|UniProtKB:P28660}; Cytoplasmic side {ECO:0000250|UniProtKB:P28660}. Note=At the interface between the lamellipodial actin meshwork and the membrane {ECO:0000250|UniProtKB:P28660}

Tissue Location

Expressed in all tissues examined except peripheral blood leukocytes, with highest expression in brain, heart, and skeletal muscle. Expressed in cells of various brain regions including Purkinje cells and dentate nucleus of the cerebellum, CA4 region and dentate gyrus of the hippocampus, and in frontal gray and white matter (PubMed:28940097).

NCKAP1 Antibody (internal region) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

NCKAP1 Antibody (internal region) - Images**NCKAP1 Antibody (internal region) - Background**

This antibody is expected to recognize both reported isoforms (NP_038464.1; NP_995314.1).

NCKAP1 Antibody (internal region) - References

Molecular cloning of a novel apoptosis-related gene, human Nap1 (NCKAP1), and its possible relation to Alzheimer disease. Suzuki T, Nishiyama K, Yamamoto A, Inazawa J, Iwaki T, Yamada T, Kanazawa I, Sakaki Y. Genomics. 2000 Jan 15;63(2):246-54. PMID: 10673335